

PATENT
DEKA:281US

APPLICATION FOR UNITED STATES LETTERS PATENT
for
PLANTS AND SEEDS OF CORN VARIETY I015036
by
THOMAS B. CARLSON

EXPRESS MAIL MAILING LABEL

NUMBER: **EL611002542US**

DATE OF DEPOSIT: **January 29, 2001**

BACKGROUND OF THE INVENTION

1. FIELD OF THE INVENTION

The present invention relates generally to the field of corn breeding. In particular, the invention relates to inbred corn seed and plants of the variety designated I015036, and
5 derivatives and tissue cultures thereof.

2. DESCRIPTION OF RELATED ART

The goal of field crop breeding is to combine various desirable traits in a single variety/hybrid. Such desirable traits include greater yield, better stalks, better roots, resistance to insecticides, herbicides, pests, and disease, tolerance to heat and drought,
10 reduced time to crop maturity, better agronomic quality, higher nutritional value, and uniformity in germination times, stand establishment, growth rate, maturity, and fruit size.

Breeding techniques take advantage of a plant's method of pollination. There are two general methods of pollination: a plant self-pollinates if pollen from one flower is transferred to the same or another flower of the same plant. A plant cross-pollinates if
15 pollen comes to it from a flower on a different plant.

Corn plants (*Zea mays L.*) can be bred by both self-pollination and cross-pollination. Both types of pollination involve the corn plant's flowers. Corn has separate male and female flowers on the same plant, located on the tassel and the ear, respectively. Natural pollination occurs in corn when wind blows pollen from the tassels to the silks that
20 protrude from the tops of the ear shoot.

Plants that have been self-pollinated and selected for type over many generations become homozygous at almost all gene loci and produce a uniform population of true breeding progeny, a homozygous plant. A cross between two such homozygous plants produces a uniform population of hybrid plants that are heterozygous for many gene loci.
25 Conversely, a cross of two plants each heterozygous at a number of loci produces a

population of hybrid plants that differ genetically and are not uniform. The resulting non-uniformity makes performance unpredictable.

The development of uniform corn plant hybrids requires the development of homozygous inbred plants, the crossing of these inbred plants, and the evaluation of the crosses. Pedigree breeding and recurrent selection are examples of breeding methods used to develop inbred plants from breeding populations. Those breeding methods combine the genetic backgrounds from two or more inbred plants or various other broad-based sources into breeding pools from which new inbred plants are developed by selfing and selection of desired phenotypes. The new inbreds are crossed with other inbred plants and the hybrids from these crosses are evaluated to determine which of those have commercial potential.

The pedigree breeding method involves crossing two genotypes. Each genotype can have one or more desirable characteristics lacking in the other; or, each genotype can complement the other. If the two original parental genotypes do not provide all of the desired characteristics, other genotypes can be included in the breeding population. Superior plants that are the products of these crosses are selfed and selected in successive generations. Each succeeding generation becomes more homogeneous as a result of self-pollination and selection. Typically, this method of breeding involves five or more generations of selfing and selection: $S_1 \rightarrow S_2$; $S_2 \rightarrow S_3$; $S_3 \rightarrow S_4$; $S_4 \rightarrow S_5$, etc. After at least five generations, the inbred plant is considered genetically pure.

Backcrossing can also be used to improve an inbred plant. Backcrossing transfers a specific desirable trait from one inbred or non-inbred source to an inbred that lacks that trait. This can be accomplished, for example, by first crossing a superior inbred (A) (recurrent parent) to a donor inbred (non-recurrent parent), which carries the appropriate locus or loci for the trait in question. The progeny of this cross are then mated back to the superior recurrent parent (A) followed by selection in the resultant progeny for the desired trait to be transferred from the non-recurrent parent. After five or more backcross generations with selection for the desired trait, the progeny are heterozygous for loci controlling the characteristic being transferred, but are like the superior parent for most or

almost all other loci. The last backcross generation would be selfed to give pure breeding progeny for the trait being transferred.

A single cross hybrid corn variety is the cross of two inbred plants, each of which has a genotype which complements the genotype of the other. The hybrid progeny of the first generation is designated F_1 . Typically, F_1 hybrids are more vigorous than their inbred parents. This hybrid vigor, or heterosis, is manifested in many polygenic traits, including markedly improved yields, better stalks, better roots, better uniformity and better insect and disease resistance. In the development of hybrids only the F_1 hybrid plants are typically sought. An F_1 single cross hybrid is produced when two inbred plants are crossed. A double cross hybrid is produced from four inbred plants crossed in pairs ($A \times B$ and $C \times D$) and then the two F_1 hybrids are crossed again $(A \times B) \times (C \times D)$.

The development of a hybrid corn variety involves three steps: (1) the selection of plants from various germplasm pools; (2) the selfing of the selected plants for several generations to produce a series of inbred plants, which, although different from each other, each breed true and are highly uniform; and (3) crossing the selected inbred plants with unrelated inbred plants to produce the hybrid progeny (F_1). During the inbreeding process in corn, the vigor of the plants decreases. Vigor is restored when two unrelated inbred plants are crossed to produce the hybrid progeny (F_1). An important consequence of the homozygosity and homogeneity of the inbred plants is that the hybrid between any two inbreds is always the same. Once the inbreds that give a superior hybrid have been identified, hybrid seed can be reproduced indefinitely as long as the homogeneity of the inbred parents is maintained. Conversely, much of the hybrid vigor exhibited by F_1 hybrids is lost in the next generation (F_2). Consequently, seed from hybrid varieties is not used for planting stock. It is not generally beneficial for farmers to save seed of F_1 hybrids. Rather, farmers purchase F_1 hybrid seed for planting every year.

North American farmers plant tens of millions of acres of corn at the present time and there are extensive national and international commercial corn breeding programs. A continuing goal of these corn breeding programs is to develop corn hybrids that are based

on stable inbred plants and have one or more desirable characteristics. To accomplish this goal, the corn breeder must select and develop superior inbred parental plants.

SUMMARY OF THE INVENTION

5 In one aspect, the present invention provides a corn plant of the variety designated I015036. Also provided are corn plants having all the physiological and morphological characteristics of the inbred corn variety I015036. The inbred corn plant of the invention may further comprise, or have, a cytoplasmic or nuclear factor that is capable of conferring male sterility or otherwise preventing self-pollination, such as by self-
10 incompatibility. Parts of the corn plant of the present invention are also provided, for example, pollen obtained from an inbred plant and an ovule of the inbred plant.

⁵⁴⁸
^{B'} The invention also concerns seed of the inbred corn variety I015036. A sample of this seed has been deposited under ATCC Accession No. - - - - -. The inbred corn seed of the invention may be provided as an essentially homogeneous population of inbred corn
15 seed of the variety designated I015036. Essentially homogeneous populations of inbred seed are those that consist essentially of the particular inbred seed, and are generally free from substantial numbers of other seed, so that the inbred seed forms between about 90% and about 100% of the total seed, and preferably, between about 95% and about 100% of the total seed. Most preferably, an essentially homogeneous population of inbred corn
20 seed will contain between about 98.5%, 99%, 99.5% and about 99.9% of inbred seed, as measured by seed grow outs. This corresponds to current commercial practice among the leading companies in the seed industry.

Therefore, in the practice of the present invention, inbred seed generally forms at least about 97% of the total seed. However, even if a population of inbred corn seed was
25 found, for some reason, to contain about 50%, or even about 20% or 15% of inbred seed, this would still be distinguished from the small fraction (generally less than 2% and preferably less than 1%) of inbred seed that may be found within a population of hybrid

seed, *e.g.*, within a commercial bag of hybrid seed. In such a bag of hybrid seed offered for sale, Federal regulations require that the hybrid seed be at least about 95% of the total seed, or be labeled as a mixture. In the most preferred practice of the invention, the female inbred seed that may be found within a bag of hybrid seed will be about 1% of the total seed, or less, and the male inbred seed that may be found within a bag of hybrid seed will be negligible, *i.e.*, will be on the order of about a maximum of 1 per 100,000, and usually less than this value.

The population of inbred corn seed of the invention can further be particularly defined as being essentially free from hybrid seed. The inbred seed population may be separately grown to provide an essentially homogeneous population of inbred corn plants designated I015036.

In another aspect of the invention, single locus converted plants of variety I015036 are provided. The single transferred locus may preferably be a dominant or recessive allele. Preferably, the single transferred locus will confer such traits as male sterility, yield stability, waxy starch, yield enhancement, industrial usage, herbicide resistance, insect resistance, resistance to bacterial, fungal, nematode or viral disease, male fertility, and enhanced nutritional quality. The single locus may be a naturally occurring maize gene introduced into the genome of the variety by backcrossing, a natural or induced mutation, or a transgene introduced through genetic transformation techniques. When introduced through transformation, a single locus may comprise one or more transgenes integrated at a single chromosomal location.

In yet another aspect of the invention, an inbred corn plant of the variety designated I015036 is provided, wherein a cytoplasmically-inherited trait has been introduced into said inbred plant. Such cytoplasmically-inherited traits are passed to progeny through the female parent in a particular cross. An exemplary cytoplasmically-inherited trait is the male sterility trait. Cytoplasmic-male sterility (CMS) is a pollen abortion phenomenon determined by the interaction between the genes in the cytoplasm and the nucleus. Alteration in the mitochondrial genome and the lack of restorer genes in the

nucleus will lead to pollen abortion. With either a normal cytoplasm or the presence of restorer gene(s) in the nucleus, the plant will produce pollen normally. A CMS plant can be pollinated by a maintainer version of the same variety, which has a normal cytoplasm but lacks the restorer gene(s) in the nucleus, and continue to be male sterile in the next
5 generation. The male fertility of a CMS plant can be restored by a restorer version of the same variety, which must have the restorer gene(s) in the nucleus. With the restorer gene(s) in the nucleus, the offspring of the male-sterile plant can produce normal pollen grains and propagate. A cytoplasmically inherited trait may be a naturally occurring maize trait or a trait introduced through genetic transformation techniques.

10 In another aspect of the invention, a tissue culture of regenerable cells of a plant of variety I015036 is provided. The tissue culture will preferably be capable of regenerating plants capable of expressing all of the physiological and morphological characteristics of the variety, and of regenerating plants having substantially the same genotype as other plants of the variety. Examples of some of the physiological and morphological
15 characteristics of the variety I015036 include characteristics related to yield, maturity, and kernel quality, each of which is specifically disclosed herein. The regenerable cells in such tissue cultures will preferably be derived from embryos, meristematic cells, immature tassels, microspores, pollen, leaves, anthers, roots, root tips, silk, flowers, kernels, ears, cobs, husks, or stalks, or from callus or protoplasts derived from those tissues. Still
20 further, the present invention provides corn plants regenerated from the tissue cultures of the invention, the plants having all the physiological and morphological characteristics of variety I015036.

In yet another aspect of the invention, processes are provided for producing corn seeds or plants, which processes generally comprise crossing a first parent corn plant with
25 a second parent corn plant, wherein at least one of the first or second parent corn plants is a plant of the variety designated I015036. These processes may be further exemplified as processes for preparing hybrid corn seed or plants, wherein a first inbred corn plant is crossed with a second corn plant of a different, distinct variety to provide a hybrid that

has, as one of its parents, the inbred corn plant variety IO15036. In these processes, crossing will result in the production of seed. The seed production occurs regardless of whether the seed is collected or not.

5 In a preferred embodiment of the invention, the first step in "crossing" comprises planting, preferably in pollinating proximity, seeds of a first and second parent corn plant, and preferably, seeds of a first inbred corn plant and a second, distinct inbred corn plant. Where the plants are not in pollinating proximity, pollination can nevertheless be accomplished by transferring a pollen or tassel bag from one plant to the other as described below.

10 A second step comprises cultivating or growing the seeds of said first and second parent corn plants into plants that bear flowers. Corn bears both male flowers (tassels) and female flowers (silks) in separate anatomical structures on the same plant.

15 A third step comprises preventing self-pollination of the plants, *i.e.*, preventing the silks of a plant from being fertilized by any plant of the same variety, including the same plant. This is preferably done by emasculating the male flowers of the first or second parent corn plant, (*i.e.*, treating or manipulating the flowers so as to prevent pollen production, in order to produce an emasculated parent corn plant). Self-incompatibility systems are also used in some hybrid crops for the same purpose. Self-incompatible plants still shed viable pollen and can pollinate plants of other varieties but are incapable of
20 pollinating themselves or other plants of the same variety.

A fourth step comprises allowing cross-pollination to occur between the first and second parent corn plants. When the plants are not in pollinating proximity, this is done by placing a bag, usually paper or glassine, over the tassels of the first plant and another bag over the silks of the incipient ear on the second plant. The bags are left in place for at least
25 24 hours. Since pollen is viable for less than 24 hours, this assures that the silks are not pollinated from other pollen sources, that any stray pollen on the tassels of the first plant is dead, and that the only pollen transferred comes from the first plant. The pollen bag over

the tassel of the first plant is then shaken vigorously to enhance release of pollen from the tassels, and the shoot bag is removed from the silks of the incipient ear on the second plant. Finally, the pollen bag is removed from the tassel of the first plant and is placed over the silks of the incipient ear of the second plant, shaken again and left in place. Yet
5 another step comprises harvesting the seeds from at least one of the parent corn plants. The harvested seed can be grown to produce a corn plant or hybrid corn plant.

The present invention also provides corn seed and plants produced by a process that comprises crossing a first parent corn plant with a second parent corn plant, wherein at least one of the first or second parent corn plants is a plant of the variety designated
10 I015036. In one embodiment of the invention, corn seed and plants produced by the process are first generation (F_1) hybrid corn seed and plants produced by crossing an inbred in accordance with the invention with another, distinct inbred. The present invention further contemplates seed of an F_1 hybrid corn plant. Therefore, certain exemplary embodiments of the invention provide an F_1 hybrid corn plant and seed thereof.
15 An example of such a hybrid which can be produced with the variety designated I015036 is the hybrid designated 8012681.

In still yet another aspect of the invention, the genetic complement of the corn plant variety designated I015036 is provided. The phrase "genetic complement" is used to refer to the aggregate of nucleotide sequences, the expression of which sequences defines
20 the phenotype of, in the present case, a corn plant, or a cell or tissue of that plant. A genetic complement thus represents the genetic make up of an inbred cell, tissue or plant, and a hybrid genetic complement represents the genetic make up of a hybrid cell, tissue or plant. The invention thus provides corn plant cells that have a genetic complement in accordance with the inbred corn plant cells disclosed herein, and plants, seeds and diploid
25 plants containing such cells.

Plant genetic complements may be assessed by genetic marker profiles, and by the expression of phenotypic traits that are characteristic of the expression of the genetic complement, *e.g.*, isozyme typing profiles. Thus, such corn plant cells may be defined as

having an SSR profile in accordance with the profile shown in Table 6, or a genetic isozyme typing profile in accordance with the profile shown in Table 7, or having both an SSR profile and an isozyme typing profile in accordance with the profiles shown in Table 6 and Table 7. It is understood that variety I015036 could also be identified by other types of genetic markers such as, for example, Simple Sequence Length Polymorphisms (SSLPs) (Williams *et al.*, 1990), Randomly Amplified Polymorphic DNAs (RAPDs), DNA Amplification Fingerprinting (DAF), Sequence Characterized Amplified Regions (SCARs), Arbitrary Primed Polymerase Chain Reaction (AP-PCR), Amplified Fragment Length Polymorphisms (AFLPs) (EP 534 858, specifically incorporated herein by reference in its entirety), and Single Nucleotide Polymorphisms (SNPs) (Wang *et al.*, 1998).

In still yet another aspect, the present invention provides hybrid genetic complements, as represented by corn plant cells, tissues, plants, and seeds, formed by the combination of a haploid genetic complement of an inbred corn plant of the invention with a haploid genetic complement of a second corn plant, preferably, another, distinct inbred corn plant. In another aspect, the present invention provides a corn plant regenerated from a tissue culture that comprises a hybrid genetic complement of this invention.

^{Sub A2} In still yet another aspect, the present invention provides a method of producing an inbred corn plant derived from the corn variety I015036, the method comprising the steps of: (a) preparing a progeny plant derived from corn variety I015036, wherein said preparing comprises crossing a plant of the corn variety I015036 with a second corn plant, and wherein a sample of the seed of corn variety I015036 has been deposited under ATCC Accession No. - - - -; (b) crossing the progeny plant with itself or a second plant to produce a seed of a progeny plant of a subsequent generation; (c) growing a progeny plant of a subsequent generation from said seed of a progeny plant of a subsequent generation and crossing the progeny plant of a subsequent generation with itself or a second plant; and (d) repeating steps (c) and (d) for an addition 3-10 generations to produce an inbred corn plant derived from the corn variety I015036. In the method, it may be desirable to

select particular plants resulting from step (c) for continued crossing according to steps (b) and (c). By selecting plants having one or more desirable traits, an inbred corn plant derived from the corn variety I015036 is obtained which possesses some of the desirable traits of corn variety I015036 as well potentially other selected traits.

5

DETAILED DESCRIPTION OF THE INVENTION

I. DEFINITIONS OF PLANT CHARACTERISTICS

Barren Plants: Plants that are barren, *i.e.*, lack an ear with grain, or have an ear with only a few scattered kernels.

Cg: *Colletotrichum graminicola* rating. Rating times 10 is approximately equal
10 to percent total plant infection.

CLN: Corn Lethal Necrosis (combination of Maize Chlorotic Mottle Virus and Maize Dwarf Mosaic virus) rating: numerical ratings are based on a severity scale where 1 = most resistant to 9 = susceptible.

Cn: *Corynebacterium nebraskense* rating. Rating times 10 is approximately equal
15 to percent total plant infection.

Cz: *Cercospora zeae-maydis* rating. Rating times 10 is approximately equal to percent total plant infection.

Dgg: *Diatraea grandiosella* girdling rating (values are percent plants girdled and stalk lodged).

Dropped Ears: Ears that have fallen from the plant to the ground.
20

Dsp: *Diabrotica* species root ratings (1 = least affected to 9 = severe pruning).

Ear-Attitude: The attitude or position of the ear at harvest scored as upright, horizontal, or pendant.

Ear-Cob Color: The color of the cob, scored as white, pink, red, or brown.

Ear-Cob Diameter: The average diameter of the cob measured at the midpoint.

5 **Ear-Cob Strength:** A measure of mechanical strength of the cobs to breakage, scored as strong or weak.

Ear-Diameter: The average diameter of the ear at its midpoint.

Ear-Dry Husk Color: The color of the husks at harvest scored as buff, red, or purple.

10 **Ear-Fresh Husk Color:** The color of the husks 1 to 2 weeks after pollination scored as green, red, or purple.

Ear-Husk Bract: The length of an average husk leaf scored as short, medium, or long.

15 **Ear-Husk Cover:** The average distance from the tip of the ear to the tip of the husks. Minimum value no less than zero.

Ear-Husk Opening: An evaluation of husk tightness at harvest scored as tight, intermediate, or open.

Ear-Length: The average length of the ear.

Ear-Number Per Stalk: The average number of ears per plant.

20 **Ear-Shank Internodes:** The average number of internodes on the ear shank.

Ear-Shank Length: The average length of the ear shank.

Ear-Shelling Percent: The average of the shelled grain weight divided by the sum of the shelled grain weight and cob weight for a single ear.

Ear-Silk Color: The color of the silk observed 2 to 3 days after silk emergence
5 scored as green-yellow, yellow, pink, red, or purple.

Ear-Taper (Shape): The taper or shape of the ear scored as conical, semi-conical, or cylindrical.

Ear-Weight: The average weight of an ear.

Early Stand: The percent of plants that emerge from the ground as determined in
10 the early spring.

ER: Ear rot rating (values approximate percent ear rotted).

Final Stand Count: The number of plants just prior to harvest.

GDUs: Growing degree units which are calculated by the Barger Method, where the heat units for a 24-h period are calculated as $GDUs = [(Maximum\ daily\ temperature + Minimum\ daily\ temperature)/2] - 50$. The highest maximum daily temperature used is 86°
15 F and the lowest minimum temperature used is 50°F.

GDUs to Shed: The number of growing degree units (GDUs) or heat units required for an inbred line or hybrid to have approximately 50% of the plants shedding pollen as measured from time of planting. GDUs to shed is determined by summing the
20 individual GDU daily values from planting date to the date of 50% pollen shed.

GDUs to Silk: The number of growing degree units for an inbred line or hybrid to have approximately 50% of the plants with silk emergence as measured from time of planting. GDUs to silk is determined by summing the individual GDU daily values from planting date to the date of 50% silking.

5 **Hc2:** *Helminthosporium carbonum* race 2 rating. Rating times 10 is approximately equal to percent total plant infection.

Hc3: *Helminthosporium carbonum* race 3 rating. Rating times 10 is approximately equal to percent total plant infection.

10 **Hm:** *Helminthosporium maydis* race 0 rating. Rating times 10 is approximately equal to percent total plant infection.

Ht1: *Helminthosporium turcicum* race 1 rating. Rating times 10 is approximately equal to percent total plant infection.

Ht2: *Helminthosporium turcicum* race 2 rating. Rating times 10 is approximately equal to percent total plant infection.

15 **HtG:** Chlorotic-lesion type resistance. += indicates the presence of Ht chlorotic-lesion type resistance; - = indicates absence of Ht chlorotic-lesion type resistance; and +/- = indicates segregation of Ht chlorotic-lesion type resistance. Rating times 10 is approximately equal to percent total plant infection.

20 **Kernel-Aleurone Color:** The color of the aleurone scored as white, pink, tan, brown, bronze, red, purple, pale purple, colorless, or variegated.

Kernel-Cap Color: The color of the kernel cap observed at dry stage, scored as white, lemon-yellow, yellow, or orange.

Kernel-Endosperm Color: The color of the endosperm scored as white, pale yellow, or yellow.

Kernel-Endosperm Type: The type of endosperm scored as normal, waxy, or opaque.

5 **Kernel-Grade:** The percent of kernels that are classified as rounds.

Kernel-Length: The average distance from the cap of the kernel to the pedicel.

Kernel-Number Per Row: The average number of kernels in a single row.

Kernel-Pericarp Color: The color of the pericarp scored as colorless, red-white crown, tan, bronze, brown, light red, cherry red, or variegated.

10 **Kernel-Row Direction:** The direction of the kernel rows on the ear scored as straight, slightly curved, spiral, or indistinct (scattered).

Kernel-Row Number: The average number of rows of kernels on a single ear.

Kernel-Side Color: The color of the kernel side observed at the dry stage, scored as white, pale yellow, yellow, orange, red, or brown.

15 **Kernel-Thickness:** The distance across the narrow side of the kernel.

Kernel-Type: The type of kernel scored as dent, flint, or intermediate.

Kernel-Weight: The average weight of a predetermined number of kernels.

Kernel-Width: The distance across the flat side of the kernel.

Kz: *Kabatiella zeae* rating. Rating times 10 is approximately equal to percent total plant infection.

Leaf-Angle: Angle of the upper leaves to the stalk scored as upright (0 to 30 degrees), intermediate (30 to 60 degrees), or lax (60 to 90 degrees).

5 **Leaf-Color:** The color of the leaves 1 to 2 weeks after pollination scored as light green, medium green, dark green, or very dark green.

Leaf-Length: The average length of the primary ear leaf.

Leaf-Longitudinal Creases: A rating of the number of longitudinal creases on the leaf surface 1 to 2 weeks after pollination. Creases are scored as absent, few, or many.

10 **Leaf-Marginal Waves:** A rating of the waviness of the leaf margin 1 to 2 weeks after pollination. Rated as none, few, or many.

Leaf-Number: The average number of leaves of a mature plant. Counting begins with the cotyledonary leaf and ends with the flag leaf.

15 **Leaf-Sheath Anthocyanin:** A rating of the level of anthocyanin in the leaf sheath 1 to 2 weeks after pollination, scored as absent, basal-weak, basal-strong, weak or strong.

Leaf-Sheath Pubescence: A rating of the pubescence of the leaf sheath. Ratings are taken 1 to 2 weeks after pollination and scored as light, medium, or heavy.

Leaf-Width: The average width of the primary ear leaf measured at its widest point.

20 **LSS:** Late season standability (values times 10 approximate percent plants lodged in disease evaluation plots).

Moisture: The moisture of the grain at harvest.

On1: *Ostrinia nubilalis* 1st brood rating (1 = resistant to 9 = susceptible).

On2: *Ostrinia nubilalis* 2nd brood rating (1 = resistant to 9 = susceptible).

Relative Maturity: A maturity rating based on regression analysis. The regression analysis is developed by utilizing check hybrids and their previously established day rating versus actual harvest moistures. Harvest moisture on the hybrid in question is determined and that moisture value is inserted into the regression equation to yield a relative maturity.

Root Lodging: Root lodging is the percentage of plants that root lodge. A plant is counted as root lodged if a portion of the plant leans from the vertical axis by approximately 30 degrees or more.

Seedling Color: Color of leaves at the 6 to 8 leaf stage.

Seedling Height: Plant height at the 6 to 8 leaf stage.

Seedling Vigor: A visual rating of the amount of vegetative growth on a 1 to 9 scale, where 1 equals best. The score is taken when the average entry in a trial is at the fifth leaf stage.

Selection Index: The selection index gives a single measure of hybrid's worth based on information from multiple traits. One of the traits that is almost always included is yield. Traits may be weighted according to the level of importance assigned to them.

Sr: *Sphacelotheca reiliana* rating is actual percent infection.

Stalk-Anthocyanin: A rating of the amount of anthocyanin pigmentation in the stalk. The stalk is rated 1 to 2 weeks after pollination as absent, basal-weak, basal-strong, weak, or strong.

Stalk-Brace Root Color: The color of the brace roots observed 1 to 2 weeks
5 after pollination as green, red, or purple.

Stalk-Diameter: The average diameter of the lowest visible internode of the stalk.

Stalk-Ear Height: The average height of the ear measured from the ground to the point of attachment of the ear shank of the top developed ear to the stalk.

Stalk-Internode Direction: The direction of the stalk internode observed after
10 pollination as straight or zigzag.

Stalk-Internode Length: The average length of the internode above the primary ear.

Stalk Lodging: The percentage of plants that did stalk lodge. Plants are counted
15 as stalk lodged if the plant is broken over or off below the ear.

Stalk-Nodes With Brace Roots: The average number of nodes having brace roots per plant.

Stalk-Plant Height: The average height of the plant as measured from the soil to the tip of the tassel.

Stalk-Tillers: The percent of plants that have tillers. A tiller is defined as a
20 secondary shoot that has developed as a tassel capable of shedding pollen.

Staygreen: Staygreen is a measure of general plant health near the time of black layer formation (physiological maturity). It is usually recorded at the time the ear husks of most entries within a trial have turned a mature color. Scoring is on a 1 to 9 basis where 1 equals best.

5 **STR:** Stalk rot rating (values represent severity rating of 1 = 25% of inoculated internode rotted to 9 = entire stalk rotted and collapsed).

10 **SVC:** Southeastern Virus Complex (combination of Maize Chlorotic Dwarf Virus and Maize Dwarf Mosaic Virus) rating; numerical ratings are based on a severity scale where 1 = most resistant to 9 = susceptible (1988 reactions are largely Maize Dwarf Mosaic Virus reactions).

Tassel-Anther Color: The color of the anthers at 50% pollen shed scored as green-yellow, yellow, pink, red, or purple.

Tassel-Attitude: The attitude of the tassel after pollination scored as open or compact.

15 **Tassel-Branch Angle:** The angle of an average tassel branch to the main stem of the tassel scored as upright (less than 30 degrees), intermediate (30 to 45 degrees), or lax (greater than 45 degrees).

Tassel-Branch Number: The average number of primary tassel branches.

20 **Tassel-Glume Band:** The closed anthocyanin band at the base of the glume scored as present or absent.

Tassel-Glume Color: The color of the glumes at 50% shed scored as green, red, or purple.

Tassel-Length: The length of the tassel measured from the base of the bottom tassel branch to the tassel tip.

Tassel-Peduncle Length: The average length of the tassel peduncle, measured from the base of the flag leaf to the base of the bottom tassel branch.

- 5 **Tassel-Pollen Shed:** A visual rating of pollen shed determined by tapping the tassel and observing the pollen flow of approximately five plants per entry. Rated on a 1 to 9 scale where 9 = sterile, 1 = most pollen.

Tassel-Spike Length: The length of the spike measured from the base of the top tassel branch to the tassel tip.

- 10 **Test Weight:** Weight of the grain in pounds for a given volume (bushel) adjusted to 15.5% moisture.

Yield: Yield of grain at harvest adjusted to 15.5% moisture.

II. OTHER DEFINITIONS

- 15 **Allele:** Any of one or more alternative forms of a gene locus, all of which alleles relate to one trait or characteristic. In a diploid cell or organism, the two alleles of a given gene occupy corresponding loci on a pair of homologous chromosomes.

- Backcrossing:** A process in which a breeder repeatedly crosses hybrid progeny back to one of the parents, for example, a first generation hybrid (F_1) with one of the parental genotypes of the F_1 hybrid.
- 20

Chromatography: A technique wherein a mixture of dissolved substances are bound to a solid support followed by passing a column of fluid across the solid support

and varying the composition of the fluid. The components of the mixture are separated by selective elution.

Crossing: The pollination of a female flower of a corn plant, thereby resulting in the production of seed from the flower.

5 **Cross-pollination:** Fertilization by the union of two gametes from different plants.

Diploid: A cell or organism having two sets of chromosomes.

10 **Electrophoresis:** A process by which particles suspended in a fluid or a gel matrix are moved under the action of an electrical field, and thereby separated according to their charge and molecular weight. This method of separation is well known to those skilled in the art and is typically applied to separating various forms of enzymes and of DNA fragments produced by restriction endonucleases.

15 **Emasculate:** The removal of plant male sex organs or the inactivation of the organs with a chemical agent or a cytoplasmic or nuclear genetic factor conferring male sterility.

Enzymes: Molecules which can act as catalysts in biological reactions.

F₁ Hybrid: The first generation progeny of the cross of two plants.

20 **Genetic Complement:** An aggregate of nucleotide sequences, the expression of which sequences defines the phenotype in corn plants, or components of plants including cells or tissue.

Genotype: The genetic constitution of a cell or organism.

Haploid: A cell or organism having one set of the two sets of chromosomes in a diploid.

Isozymes: Detectable variants of an enzyme, the variants catalyzing the same reaction(s) but differing from each other, *e.g.*, in primary structure and/or electrophoretic mobility. The differences between isozymes are under single gene, codominant control. Consequently, electrophoretic separation to produce band patterns can be equated to different alleles at the DNA level. Structural differences that do not alter charge cannot be detected by this method.

Isozyme typing profile: A profile of band patterns of isozymes separated by electrophoresis that can be equated to different alleles at the DNA level.

Linkage: A phenomenon wherein alleles on the same chromosome tend to segregate together more often than expected by chance if their transmission was independent.

Marker: A readily detectable phenotype, preferably inherited in codominant fashion (both alleles at a locus in a diploid heterozygote are readily detectable), with no environmental variance component, *i.e.*, heritability of 1.

Sub B37 1015036: The corn plant variety from which seeds having ATCC Accession No. - - - - were obtained, as well as plants grown from those seeds.

Phenotype: The detectable characteristics of a cell or organism, which characteristics are the manifestation of gene expression.

Quantitative Trait Loci (QTL): Genetic loci that contribute, at least in part, certain numerically representable traits that are usually continuously distributed.

Regeneration: The development of a plant from tissue culture.

SSR profile: A profile of simple sequence repeats used as genetic markers and scored by gel electrophoresis following PCR™ amplification using flanking oligonucleotide primers.

5 **Self-pollination:** The transfer of pollen from the anther to the stigma of the same plant.

10 **Single Locus Converted (Conversion) Plant:** Plants which are developed by a plant breeding technique called backcrossing wherein essentially all of the desired morphological and physiological characteristics of an inbred are recovered in addition to the characteristics conferred by the single locus transferred into the inbred *via* the backcrossing technique. A single locus may comprise one gene, or in the case of transgenic plants, one or more transgenes integrated into the host genome at a single site (locus).

Tissue Culture: A composition comprising isolated cells of the same or a different type or a collection of such cells organized into parts of a plant.

15 **Transgene:** A genetic sequence which has been introduced into the nuclear or chloroplast genome of a maize plant by a genetic transformation technique.

20 The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

25

III. INBRED CORN PLANT I015036

In accordance with one aspect of the present invention, there is provided a novel inbred corn plant variety designated I015036. Inbred corn plant I015036 can be compared to inbred corn plants FBLL and FBMU. I015036 differs significantly (at the 1%, 5%, or 5 10% level) from these inbred lines in several aspects (Table 1 and Table 2).

Table 1: Comparison of I015036 with FBLL

	I015036	FBLL	DIFF	P VALUE
BARREN %	2.8	3.0	-0.2	0.556
EHT INCH	30.5	29.9	0.6	0.872
FINAL	64.8	60.8	4.0	0.005*
MST %	17.7	22.8	-5.1	0.043+
PHT INCH	77.9	70.3	7.6	0.120
RTL %	1.6	0.5	1.1	0.000**
SHED GDU	1394.4	1472.8	-78.4	0.950
SILK GDU	1405.5	1491.8	-86.3	0.572
STL %	2.1	3.0	-0.9	0.007*
YLD BU/A	83.8	66.4	17.4	0.177

*Significance Levels are indicated as: + = 10%, * = 5%, ** = 1%.*

- 10 Legend Abbreviations:
BARREN % = Barren Plants (percent)
EHT INCH = Ear Height (inches)
FINAL = Final Stand
MST % = Moisture (percent)

- PHT INCH = Plant Height (inches)
 RTL % = Root Lodging (percent)
 SHED GDU = GDUs to Shed
 SILK GDU = GDUs to Silk
 5 STL % = Stalk Lodging (percent)
 YLD BU/A = Yield (bushels/acre)

Table 2: Comparison of I015036 with FBMU

10

	I015036	FBMU	DIFF	P VALUE
BARREN %	2.8	2.6	0.2	0.998
EHT INCH	30.5	27.7	2.8	0.876
FINAL	64.8	59.1	5.7	0.005*
MST %	17.7	18.9	-1.2	0.655
PHT INCH	77.9	65.6	12.3	0.055+
RTL %	1.6	4.8	-3.2	0.000**
SHED GDU	1394.4	1398.7	-4.3	0.680
SILK GDU	1405.5	1400.6	4.9	0.799
STL %	2.1	6.2	-4.1	0.000**
YLD BU/A	83.8	78.7	5.1	0.396

*Significance Levels are indicated as: + = 10%, * = 5%, ** = 1%.*

Legend Abbreviations:

- BARREN % = Barren Plants (percent)
 15 EHT INCH = Ear Height (inches)
 FINAL = Final Stand
 MST % = Moisture (percent)
 PHT INCH = Plant Height (inches)
 RTL % = Root Lodging (percent)

SHED GDU = GDUs to Shed
 SILK GDU = GDUs to Silk
 STL % = Stalk Lodging (percent)
 YLD BU/A =Yield(bushels/acre)

5

A. Origin and Breeding History

Inbred plant I015036 was derived from a cross between the lines FBMU and FBLL. The origin and breeding history of Corn Variety I015036 can be summarized as follows:

10

Winter 1991 The inbred line FBMU (a proprietary DEKALB Genetics Corporation Inbred) was crossed to the inbred line FBLL (a proprietary DEKALB Genetics Corporation Inbred) in nursery rows E77 x E76.

15

Winter 1991/92 The F1 seed was grown and self-pollinated in nursery row 83-176.

Summer 1993 The F2 seed was grown and self-pollinated in nursery rows 42-25 through 42-1.

Winter 1993/94 F3 Ears were grown ear-to-row and self-pollinated. F4 ears were selected in nursery rows 12C - 134.

20

Summer 1994 F4 ears were grown ear-to-row and self-pollinated. F5 ears were selected in nursery rows 281-8.

Summer 1995 F5 ears were grown ear-to-row and self-pollinated. F6 ears from nursery row 295-27 were selected.

25

Winter 1995/96 F6 ears were grown ear-to-row and self-pollinated. F7 ears from nursery row 6J 33-34 were selected.

Summer 1996 F7 ears were grown ear-to-row and self-pollinated. F8 ears were selected from nursery row 384-15.

Winter 1996/97 F8 ears were grown ear-to-row and self-pollinated. Final selection was completed in nursery rows 6J16 - 44 - 43.

30

Corn variety I015036 shows uniformity and stability within the limits of environmental influence for the traits described hereinafter in Table 3. I015036 has been

self-pollinated and ear-rowed a sufficient number of generations with careful attention paid to uniformity of plant type to ensure homozygosity and phenotypic stability. No variant traits have been observed or are expected in I015036.

Inbred corn plants can be reproduced by planting the seeds of the inbred corn plant I015036, growing the resulting corn plants under self-pollinating or sib-pollinating conditions with adequate isolation using standard techniques well known to an artisan skilled in the agricultural arts. Seeds can be harvested from such a plant using standard, well known procedures.

B. Phenotypic Description

In accordance with another aspect of the present invention, there is provided a corn plant having the physiological and morphological characteristics of corn plant I015036. A description of the physiological and morphological characteristics of corn plant I015036 is presented in Table 3.

Table 3: Morphological Traits for the I015036 Phenotype

CHARACTERISTIC		VALUE		
		I015036	FBL	FBM
1.	STALK			
	Diameter (width) cm.	2.2	2.2	2.5
	Anthocyanin	Absent	Absent	Weak
	Brace Root Color	Dark	Dark	-
	Nodes With Brace Roots	1.1	2.3	1.5
	Internode Direction	Straight	Straight	Straight
	Internode Length cm.	13.7	13.9	13.0
2.	LEAF			
	Color	Dark Green	Green	Green
	Length cm.	72.3	61.2	18.0
	Width cm.	9.0	7.1	2.2

CHARACTERISTIC		VALUE		
		I015036	FBLL	FBMU
	Sheath Anthocyanin	Basal-Strong	Weak	Weak
	Sheath Pubescence	Moderate	Moderate	Light
	Marginal Waves	Few	Few	Few
	Longitudinal Creases	Moderate	Moderate	Few

3.	TASSEL			
	Length cm.	35.9	33.5	30.7
	Spike Length cm.	18.1	16.7	18.4
	Peduncle Length cm.	10.2	9.8	9.1
	Branch Number	11.7	20.6	25.7
	Anther Color	Pink	Pink	Red
	Glume Color	Green	Green	Green
	Glume Band	Absent	Absent	Absent

4.	EAR			
	Silk Color	Green-Yellow	Green-Yellow	Pink
	Number Per Stalk	1.0	1.0	1.3
	Position (attitude)	Upright	Pendant	Upright
	Length cm.	16.6	14.5	17.5
	Shape	Semi-Conical	Semi-Conical	Semi-Conical
	Diameter cm.	4.1	4.1	4.1
	Weight gm.	161.1	115.7	120.3
	Shank Length cm.	9.2	8.0	14.2
	Husk Bract	Short	Short	Short
	Husk Cover cm.	3.8	5.5	3.6
	Husk Opening	Loose	Intermediate	Intermediate
	Husk Color Fresh	Green	Green	Green
	Husk Color Dry	Buff	Buff	Buff
	Cob Diameter cm.	2.3	2.5	2.3
	Cob Color	Red	Red	Red
	Shelling Percent	85.7	83.9	83.8

5.	KERNEL			
	Row Number	14.7	17.4	13.0
	Number Per Row	36.8	29.5	33.1
	Row Direction	Straight	Slightly Curved	Slightly Curved
	Type	Dent	Dent	Dent
	Cap Color	Orange	Yellow	Yellow
	Side Color	Orange	Deep-Yellow	Deep-Yellow
	Length (depth) mm.	10.7	11.1	10.8
	Width mm.	7.6	6.7	8.7

CHARACTERISTIC		VALUE		
		I015036	FBLL	FBMU
	Thickness	3.8	4.0	4.7
	Weight of 1000K gm.	273.0	228.1	308.6
	Endosperm Type	Normal	Normal	Normal
	Endosperm Color	Yellow	Yellow	Yellow

**These are typical values. Values may vary due to environment. Other values that are substantially equivalent are also within the scope of the invention. Substantially equivalent refers to quantitative traits that when compared do not show statistical differences of their means.*

5

C. DEPOSIT INFORMATION

Sub 104 > A representative deposit of 2500 seeds of the inbred corn variety designated I015036 has been made with the American Type Culture Collection (ATCC), 10801 University Blvd., Manassas, VA on (_____, ____). Those deposited seeds have been assigned ATCC Accession No. - - - - -. The deposit was made in accordance with the terms and provisions of the Budapest Treaty relating to deposit of microorganisms and was made for a term of at least thirty (30) years and at least five (05) years after the most recent request for the furnishing of a sample of the deposit is received by the depository, or for the effective term of the patent, whichever is longer, and will be replaced if it becomes non-viable during that period.

IV. SINGLE LOCUS CONVERSIONS

When the term inbred corn plant is used in the context of the present invention, this also includes any single locus conversions of that inbred. The term single locus converted plant as used herein refers to those corn plants which are developed by a plant breeding technique called backcrossing wherein essentially all of the desired morphological and physiological characteristics of an inbred are recovered in addition to the single locus transferred into the inbred *via* the backcrossing technique. Backcrossing methods can be used with the present invention to improve or introduce a characteristic into the inbred.

The term backcrossing as used herein refers to the repeated crossing of a hybrid progeny back to one of the parental corn plants for that inbred. The parental corn plant which contributes the locus or loci for the desired characteristic is termed the nonrecurrent or donor parent. This terminology refers to the fact that the nonrecurrent parent is used one
5 time in the backcross protocol and therefore does not recur. The parental corn plant to which the locus or loci from the nonrecurrent parent are transferred is known as the recurrent parent as it is used for several rounds in the backcrossing protocol (Poehlman *et al.*, 1995; Fehr, 1987; Sprague and Dudley, 1988). In a typical backcross protocol, the original inbred of interest (recurrent parent) is crossed to a second inbred (nonrecurrent
10 parent) that carries the single locus of interest to be transferred. The resulting progeny from this cross are then crossed again to the recurrent parent and the process is repeated until a corn plant is obtained wherein essentially all of the desired morphological and physiological characteristics of the recurrent parent are recovered in the converted plant, in addition to the single transferred locus from the nonrecurrent parent. The backcross
15 process may be accelerated by the use of genetic markers, such as SSR, RFLP, SNP or AFLP markers to identify plants with the greatest genetic complement from the recurrent parent.

The selection of a suitable recurrent parent is an important step for a successful backcrossing procedure. The goal of a backcross protocol is to alter or substitute a single
20 trait or characteristic in the original inbred. To accomplish this, a single locus of the recurrent inbred is modified or substituted with the desired locus from the nonrecurrent parent, while retaining essentially all of the rest of the desired genetic, and therefore the desired physiological and morphological constitution of the original inbred. The choice of the particular nonrecurrent parent will depend on the purpose of the backcross; one of the
25 major purposes is to add some commercially desirable, agronomically important trait to the plant. The exact backcrossing protocol will depend on the characteristic or trait being altered to determine an appropriate testing protocol. Although backcrossing methods are simplified when the characteristic being transferred is a dominant allele, a recessive allele

may also be transferred. In this instance it may be necessary to introduce a test of the progeny to determine if the desired characteristic has been successfully transferred.

Many single locus traits have been identified that are not regularly selected for in the development of a new inbred but that can be improved by backcrossing techniques.

5 Single locus traits may or may not be transgenic; examples of these traits include, but are not limited to, male sterility, waxy starch, herbicide resistance, resistance for bacterial, fungal, or viral disease, insect resistance, male fertility, enhanced nutritional quality, industrial usage, yield stability, and yield enhancement. These genes are generally inherited through the nucleus, but may be inherited through the cytoplasm. Some known

10 exceptions to this are genes for male sterility, some of which are inherited cytoplasmically, but still act as single locus traits. A number of exemplary single locus traits are described in, for example, PCT Application WO 95/06128, the disclosure of which is specifically incorporated herein by reference.

Examples of genes conferring male sterility include those disclosed in U.S. Patent

15 No. 3,861,709, U.S. Patent No. 3,710,511, U.S. Patent No. 4,654,465, U.S. Patent No. 5,625,132, and U.S. Patent No. 4,727,219, each of the disclosures of which are specifically incorporated herein by reference in their entirety. A particularly useful type of male sterility gene is one which can be induced by exposure to a chemical agent, for example, a herbicide (U.S. Patent Ser. No. 08/927,368, filed September 11, 1997, the

20 disclosure of which is specifically incorporated herein by reference in its entirety). Both inducible and non-inducible male sterility genes can increase the efficiency with which hybrids are made, in that they eliminate the need to physically emasculate the corn plant used as a female in a given cross.

Where one desires to employ male-sterility systems with a corn plant in accordance

25 with the invention, it may be beneficial to also utilize one or more male-fertility restorer genes. For example, where cytoplasmic male sterility (CMS) is used, hybrid seed production requires three inbred lines: (1) a cytoplasmically male-sterile line having a CMS cytoplasm; (2) a fertile inbred with normal cytoplasm, which is isogenic with the

CMS line for nuclear genes ("maintainer line"); and (3) a distinct, fertile inbred with normal cytoplasm, carrying a fertility restoring gene ("restorer" line). The CMS line is propagated by pollination with the maintainer line, with all of the progeny being male sterile, as the CMS cytoplasm is derived from the female parent. These male sterile plants
5 can then be efficiently employed as the female parent in hybrid crosses with the restorer line, without the need for physical emasculation of the male reproductive parts of the female parent.

The presence of a male-fertility restorer gene results in the production of fully fertile F₁ hybrid progeny. If no restorer gene is present in the male parent, male-sterile
10 hybrids are obtained. Such hybrids are useful where the vegetative tissue of the corn plant is utilized, *e.g.*, for silage, but in most cases, the seeds will be deemed the most valuable portion of the crop, so fertility of the hybrids in these crops must be restored. Therefore, one aspect of the current invention concerns the inbred corn plant I015036 comprising a single gene capable of restoring male fertility in an otherwise male-sterile inbred or hybrid
15 plant. Examples of male-sterility genes and corresponding restorers which could be employed with the inbred of the invention are well known to those of skill in the art of plant breeding and are disclosed in, for instance, U.S. Patent No. 5,530,191; U.S. Patent No. 5,689,041; U.S. Patent No. 5,741,684; and U.S. Patent No. 5,684,242, the disclosures of which are each specifically incorporated herein by reference in their entirety.

20 Direct selection may be applied where a single locus acts as a dominant trait. An example of a dominant trait is the herbicide resistance trait. For this selection process, the progeny of the initial cross are sprayed with the herbicide prior to the backcrossing. The spraying eliminates any plants which do not have the desired herbicide resistance characteristic, and only those plants which have the herbicide resistance gene are used in
25 the subsequent backcross. This process is then repeated for all additional backcross generations.

Many useful single locus traits are those which are introduced by genetic transformation techniques. Methods for the genetic transformation of maize are known to

those of skill in the art. For example, methods which have been described for the genetic transformation of maize include electroporation (U.S. Patent No. 5,384,253), electrotransformation (U.S. Patent No. 5,371,003), microprojectile bombardment (U.S. Patent No. 5,550,318; U.S. Patent No. 5,736,369, U.S. Patent No. 5,538,880; and PCT Publication WO 95/06128), *Agrobacterium*-mediated transformation (U.S. Patent No. 5,591,616 and E.P. Publication EP672752), direct DNA uptake transformation of protoplasts (Omirulleh *et al.*, 1993) and silicon carbide fiber-mediated transformation (U.S. Patent No. 5,302,532 and U.S. Patent No. 5,464,765).

A type of single locus trait which can be introduced by genetic transformation (U.S. Patent No. 5,554,798) and has particular utility is a gene which confers resistance to the herbicide glyphosate. Glyphosate inhibits the action of the enzyme EPSPS, which is active in the biosynthetic pathway of aromatic amino acids. Inhibition of this enzyme leads to starvation for the amino acids phenylalanine, tyrosine, and tryptophan and secondary metabolites derived therefrom. Mutants of this enzyme are available which are resistant to glyphosate. For example, U.S. Patent 4,535,060 describes the isolation of EPSPS mutations which confer glyphosate resistance upon organisms having the *Salmonella typhimurium* gene for EPSPS, *aroA*. A mutant EPSPS gene having similar mutations has also been cloned from *Zea mays*. The mutant gene encodes a protein with amino acid changes at residues 102 and 106 (PCT Publication WO 97/04103). When a plant comprises such a gene, a herbicide resistant phenotype results.

Plants having inherited a transgene comprising a mutated EPSPS gene may, therefore, be directly treated with the herbicide glyphosate without the result of significant damage to the plant. This phenotype provides farmers with the benefit of controlling weed growth in a field of plants having the herbicide resistance trait by application of the broad spectrum herbicide glyphosate. For example, one could apply the herbicide ROUNDUP™, a commercial formulation of glyphosate manufactured and sold by the Monsanto Company, over the top in fields where glyphosate resistant corn plants are grown. The herbicide application rates may typically range from 4 ounces of ROUNDUP™ to 256 ounces ROUNDUP™ per acre. More preferably, about 16 ounces

to about 64 ounces per acre of ROUNDUP™ may be applied to the field. However, the application rate may be increased or decreased as needed, based on the abundance and / or type of weeds being treated. Additionally, depending on the location of the field and weather conditions, which will influence weed growth and the type of weed infestation, it may be desirable to conduct further glyphosate treatments. The second glyphosate application will also typically comprise an application rate of about 16 ounces to about 64 ounces of ROUNDUP™ per acre treated. Again, the treatment rate may be adjusted based on field conditions. Such methods of application of herbicides to agricultural crops are well known in the art and are summarized in general in Anderson (1983).

10 Alternatively, more than one single locus trait may be introgressed into an elite inbred by the method of backcross conversion. A selectable marker gene and a gene encoding a protein which confers a trait of interest may be simultaneously introduced into a maize plant as a result of genetic transformation. Usually one or more introduced genes will integrate into a single chromosome site in the host cell's genome. For example, a
15 selectable marker gene encoding phosphinothricin acetyl transferase (PPT) (*e.g.*, a bar gene) and conferring resistance to the active ingredient in some herbicides by inhibiting glutamine synthetase, and a gene encoding an endotoxin from *Bacillus thuringiensis* (Bt) and conferring resistance to particular classes of insects, *e.g.*, lepidopteran insects, in particular the European Corn Borer, may be simultaneously introduced into a host
20 genome. Furthermore, through the process of backcross conversion more than one transgenic trait may be transferred into an elite inbred.

The waxy characteristic is an example of a recessive trait. In this example, the progeny resulting from the first backcross generation (BC1) must be grown and selfed. A test is then run on the selfed seed from the BC1 plant to determine which BC1 plants
25 carried the recessive gene for the waxy trait. In other recessive traits additional progeny testing, for example growing additional generations such as the BC1S1, may be required to determine which plants carry the recessive gene.

V. ORIGIN AND BREEDING HISTORY OF AN EXEMPLARY SINGLE LOCUS CONVERTED PLANT

85DGD1 MLms is a single locus conversion of 85DGD1 to cytoplasmic male sterility. 85DGD1 MLms was derived using backcross methods. 85DGD1 (a proprietary inbred of DEKALB Genetics Corporation) was used as the recurrent parent and MLms, a germplasm source carrying ML cytoplasmic sterility, was used as the nonrecurrent parent. The breeding history of the single locus converted inbred 85DGD1 MLms can be summarized as follows:

Hawaii Nurseries Planting Date 04-02-1992	Made up S-O: Female row 585 male row 500
Hawaii Nurseries Planting Date 07-15-1992	S-O was grown and plants were backcrossed times 85DGD1 (rows 444 ' 443)
Hawaii Nurseries Planting Date 11-18-1992	Bulked seed of the BC1 was grown and backcrossed times 85DGD1 (rows V3-27 ' V3-26)
Hawaii Nurseries Planting Date 04-02-1993	Bulked seed of the BC2 was grown and backcrossed times 85DGD1 (rows 37 ' 36)
Hawaii Nurseries Planting Date 07-14-1993	Bulked seed of the BC3 was grown and backcrossed times 85DGD1 (rows 99 ' 98)
Hawaii Nurseries Planting Date 10-28-1993	Bulked seed of BC4 was grown and backcrossed times 85DGD1 (rows KS-63 ' KS-62)
Summer 1994	A single ear of the BC5 was grown and backcrossed times 85DGD1 (MC94-822 ' MC94-822-7)

Winter 1994

Bulked seed of the BC6 was grown and backcrossed times 85DGD1 (3Q-1 ' 3Q-2)

Summer 1995

Seed of the BC7 was bulked and named 85DGD1 MLms.

VI. TISSUE CULTURES AND *IN VITRO* REGENERATION OF CORN PLANTS

A further aspect of the invention relates to tissue cultures of the corn plant designated I015036. As used herein, the term "tissue culture" indicates a composition comprising isolated cells of the same or a different type or a collection of such cells organized into parts of a plant. Exemplary types of tissue cultures are protoplasts, calli and plant cells that are intact in plants or parts of plants, such as embryos, pollen, flowers, kernels, ears, cobs, leaves, husks, stalks, roots, root tips, anthers, silk, and the like. In a preferred embodiment, the tissue culture comprises embryos, protoplasts, meristematic cells, pollen, leaves or anthers derived from immature tissues of these plant parts. Means for preparing and maintaining plant tissue cultures are well known in the art (U.S. Patent No. 5,538,880; and U.S. Patent No. 5,550,318, each incorporated herein by reference in their entirety). By way of example, a tissue culture comprising organs such as tassels or anthers has been used to produce regenerated plants (U.S. Patent No. 5,445,961 and U.S. Patent No. 5,322,789; the disclosures of which are incorporated herein by reference).

VII. TASSEL/ANTHER CULTURE

Tassels contain anthers which in turn enclose microspores. Microspores develop into pollen. For anther/microspore culture, if tassels are the plant composition, they are preferably selected at a stage when the microspores are uninucleate, that is, include only one, rather than 2 or 3 nuclei. Methods to determine the correct stage are well known to those skilled in the art and include mitramycin fluorescent staining (Pace *et al.*, 1987),

trypan blue (preferred) and acetocarmine squashing. The mid-uninucleate microspore stage has been found to be the developmental stage most responsive to the subsequent methods disclosed to ultimately produce plants.

Although microspore-containing plant organs such as tassels can generally be
5 pretreated at any cold temperature below about 25°C, a range of 4 to 25°C is preferred, and a range of 8 to 14°C is particularly preferred. Although other temperatures yield embryoids and regenerated plants, cold temperatures produce optimum response rates compared to pretreatment at temperatures outside the preferred range. Response rate is measured as either the number of embryoids or the number of regenerated plants per
10 number of microspores initiated in culture. Exemplary methods of microspore culture are disclosed in, for example, U.S. Patent No. 5,322,789 and U.S. Patent No 5,445,961, the disclosures of which are specifically incorporated herein by reference.

Although not required, when tassels are employed as the plant organ, it is generally preferred to sterilize their surface. Following surface sterilization of the tassels, for
15 example, with a solution of calcium hypochloride, the anthers are removed from about 70 to 150 spikelets (small portions of the tassels) and placed in a preculture or pretreatment medium. Larger or smaller amounts can be used depending on the number of anthers.

When one elects to employ tassels directly, tassels are preferably pretreated at a cold temperature for a predefined time, preferably at 10°C for about 4 days. After
20 pretreatment of a whole tassel at a cold temperature, dissected anthers are further pretreated in an environment that diverts microspores from their developmental pathway. The function of the preculture medium is to switch the developmental program from one of pollen development to that of embryoid/callus development. An embodiment of such an environment in the form of a preculture medium includes a sugar alcohol, for example
25 mannitol or sorbitol, inositol or the like. An exemplary synergistic combination is the use of mannitol at a temperature of about 10°C for a period ranging from about 10 to 14 days. In a preferred embodiment, 3 ml of 0.3 M mannitol combined with 50 mg/l of ascorbic acid, silver nitrate, and colchicine is used for incubation of anthers at 10°C for between 10

and 14 days. Another embodiment is to substitute sorbitol for mannitol. The colchicine produces chromosome doubling at this early stage. The chromosome doubling agent is preferably only present at the preculture stage.

5 It is believed that the mannitol or other similar carbon structure or environmental stress induces starvation and functions to force microspores to focus their energies on entering developmental stages. The cells are unable to use, for example, mannitol as a carbon source at this stage. It is believed that these treatments confuse the cells causing them to develop as embryoids and plants from microspores. Dramatic increases in development from these haploid cells, as high as 25 embryoids in 10^4 microspores, have
10 resulted from using these methods.

In embodiments where microspores are obtained from anthers, microspores can be released from the anthers into an isolation medium following the mannitol preculture step. One method of release is by disruption of the anthers, for example, by chopping the anthers into pieces with a sharp instrument, such as a razor blade, scalpel, or Waring
15 blender. The resulting mixture of released microspores, anther fragments, and isolation medium are then passed through a filter to separate microspores from anther wall fragments. An embodiment of a filter is a mesh, more specifically, a nylon mesh of about 112 μ m pore size. The filtrate which results from filtering the microspore-containing solution is preferably relatively free of anther fragments, cell walls, and other debris.

20 In a preferred embodiment, isolation of microspores is accomplished at a temperature below about 25°C and preferably, at a temperature of less than about 15°C. Preferably, the isolation media, dispersing tool (e.g., razor blade), funnels, centrifuge tubes, and dispersing container (e.g., petri dish) are all maintained at the reduced temperature during isolation. The use of a precooled dispersing tool to isolate maize
25 microspores has been reported (Gaillard *et al.*, 1991).

Where appropriate and desired, the anther filtrate is then washed several times in isolation medium. The purpose of the washing and centrifugation is to eliminate any toxic

compounds which are contained in the non-microspore part of the filtrate and are created by the chopping process. The centrifugation is usually done at decreasing spin speeds, for example, 1000, 750, and finally 500 rpms. The result of the foregoing steps is the preparation of a relatively pure tissue culture suspension of microspores that are relatively
5 free of debris and anther remnants.

To isolate microspores, an isolation media is preferred. An isolation media is used to separate microspores from the anther walls while maintaining their viability and embryogenic potential. An illustrative embodiment of an isolation media includes a 6% sucrose or maltose solution combined with an antioxidant such as 50 mg/l of ascorbic
10 acid, 0.1 mg/l biotin, and 400 mg/l of proline, combined with 10 mg/l of nicotinic acid and 0.5 mg/l AgNO_3 . In another embodiment, the biotin and proline are omitted.

An isolation media preferably has a higher antioxidant level where it is used to isolate microspores from a donor plant (a plant from which a plant composition containing a microspore is obtained) that is field grown in contrast to greenhouse grown. A preferred
15 level of ascorbic acid in an isolation medium is from about 50 mg/l to about 125 mg/l and, more preferably, from about 50 mg/l to about 100 mg/l.

One can find particular benefit in employing a support for the microspores during culturing and subculturing. Any support that maintains the cells near the surface can be used. The microspore suspension is layered onto a support, for example by pipetting.
20 There are several types of supports which are suitable and are within the scope of the invention. An illustrative embodiment of a solid support is a TRANSWELL® culture dish. Another embodiment of a solid support for development of the microspores is a bilayer plate wherein liquid media is on top of a solid base. Other embodiments include a mesh or a millipore filter. Preferably, a solid support is a nylon mesh in the shape of a raft.
25 A raft is defined as an approximately circular support material which is capable of floating slightly above the bottom of a tissue culture vessel, for example, a petri dish, of about a 60 or 100 mm size, although any other laboratory tissue culture vessel will suffice. In an illustrative embodiment, a raft is about 55 mm in diameter.

Culturing isolated microspores on a solid support, for example, on a 10 mm pore nylon raft floating on 2.2 ml of medium in a 60 mm petri dish, prevents microspores from sinking into the liquid medium and thus avoiding low oxygen tension. These types of cell supports enable the serial transfer of the nylon raft with its associated microspore/embryoids ultimately to full strength medium containing activated charcoal and solidified with, for example, GELRITE™ (solidifying agent). The charcoal is believed to absorb toxic wastes and intermediaries. The solid medium allows embryoids to mature.

The liquid medium passes through the mesh while the microspores are retained and supported at the medium-air interface. The surface tension of the liquid medium in the petri dish causes the raft to float. The liquid is able to pass through the mesh; consequently, the microspores stay on top. The mesh remains on top of the total volume of liquid medium. An advantage of the raft is to permit diffusion of nutrients to the microspores. Use of a raft also permits transfer of the microspores from dish to dish during subsequent subculture with minimal loss, disruption, or disturbance of the induced embryoids that are developing. The rafts represent an advantage over the multi-welled TRANSWELL® plates, which are commercially available from COSTAR, in that the commercial plates are expensive. Another disadvantage of these plates is that to achieve the serial transfer of microspores to subsequent media, the membrane support with cells must be peeled off the insert in the wells. This procedure does not produce as good a yield nor as efficient transfers, as when a mesh is used as a vehicle for cell transfer.

The culture vessels can be further defined as either (1) a bilayer 60 mm petri plate wherein the bottom 2 ml of medium are solidified with 0.7% agarose overlaid with 1 mm of liquid containing the microspores; (2) a nylon mesh raft wherein a wafer of nylon is floated on 1.2 ml of medium and 1 ml of isolated microspores is pipetted on top; or (3) TRANSWELL® plates wherein isolated microspores are pipetted onto membrane inserts which support the microspores at the surface of 2 ml of medium.

After the microspores have been isolated, they are cultured in a low strength anther culture medium until about the 50 cell stage when they are subcultured onto an

embryoid/callus maturation medium. Medium is defined at this stage as any combination of nutrients that permit the microspores to develop into embryoids or callus. Many examples of suitable embryoid/callus promoting media are well known to those skilled in the art. These media will typically comprise mineral salts, a carbon source, vitamins, and growth regulators. A solidifying agent is optional. A preferred embodiment of such a media is referred to as "D medium," which typically includes 6N1 salts, AgNO₃ and sucrose or maltose.

In an illustrative embodiment, 1 ml of isolated microspores are pipetted onto a 10 mm nylon raft and the raft is floated on 1.2 ml of medium "D," containing sucrose or preferably maltose. Both calli and embryoids can develop. Calli are undifferentiated aggregates of cells. Type I is a relatively compact, organized, and slow growing callus. Type II is a soft, friable, and fast-growing one. Embryoids are aggregates exhibiting some embryo-like structures. The embryoids are preferred for subsequent steps to regenerating plants. Culture medium "D" is an embodiment of medium that follows the isolation medium and replaces it. Medium "D" promotes growth to an embryoid/callus. This medium comprises 6N1 salts at 1/8 the strength of a basic stock solution (major components) and minor components, plus 12% sucrose, or preferably 12% maltose, 0.1 mg/l B1, 0.5 mg/l nicotinic acid, 400 mg/l proline and 0.5 mg/l silver nitrate. Silver nitrate is believed to act as an inhibitor to the action of ethylene. Multi-cellular structures of approximately 50 cells each generally arise during a period of 12 days to 3 weeks. Serial transfer after a two week incubation period is preferred.

After the petri dish has been incubated for an appropriate period of time, preferably two weeks in the dark at a predefined temperature, a raft bearing the dividing microspores is transferred serially to solid based media which promote embryo maturation. In an illustrative embodiment, the incubation temperature is 30°C and the mesh raft supporting the embryoids is transferred to a 100 mm petri dish containing the 6N1-TGR-4P medium, an "anther culture medium." This medium contains 6N1 salts, supplemented with 0.1 mg/l TIBA, 12% sugar (sucrose, maltose, or a combination thereof), 0.5%

activated charcoal, 400 mg/l proline, 0.5 mg/l B, 0.5 mg/l nicotinic acid, and 0.2 percent GELRITE™ (solidifying agent) and is capable of promoting the maturation of the embryoids. Higher quality embryoids, that is, embryoids which exhibit more organized development, such as better shoot meristem formation without precocious germination, were typically obtained with the transfer to full strength medium compared to those resulting from continuous culture using only, for example, the isolated microspore culture (IMC) Medium "D." The maturation process permits the pollen embryoids to develop further in route toward the eventual regeneration of plants. Serial transfer occurs to full strength solidified 6N1 medium using either the nylon raft, the TRANSWELL® membrane, or bilayer plates, each one requiring the movement of developing embryoids to permit further development into physiologically more mature structures. In an especially preferred embodiment, microspores are isolated in an isolation media comprising about 6% maltose, cultured for about two weeks in an embryoid/calli induction medium comprising about 12% maltose and then transferred to a solid medium comprising about 12% sucrose.

At the point of transfer of the raft, after about two weeks of incubation, embryoids exist on a nylon support. The purpose of transferring the raft with the embryoids to a solidified medium after the incubation is to facilitate embryo maturation. Mature embryoids at this point are selected by visual inspection indicated by zygotic embryo-like dimensions and structures and are transferred to the shoot initiation medium. It is preferred that shoots develop before roots, or that shoots and roots develop concurrently. If roots develop before shoots, plant regeneration can be impaired. To produce solidified media, the bottom of a petri dish of approximately 100 mm is covered with about 30 ml of 0.2% GELRITE™ solidified medium. A sequence of regeneration media are used for whole plant formation from the embryoids.

During the regeneration process, individual embryoids are induced to form plantlets. The number of different media in the sequence can vary depending on the specific protocol used. Finally, a rooting medium is used as a prelude to transplanting to

soil. When plantlets reach a height of about 5 cm, they are then transferred to pots for further growth into flowering plants in a greenhouse by methods well known to those skilled in the art.

Plants have been produced from isolated microspore cultures by the methods disclosed herein, including self-pollinated plants. The rate of embryoid induction was much higher with the synergistic preculture treatment consisting of a combination of stress factors, including a carbon source which can be capable of inducing starvation, a cold temperature, and colchicine, than has previously been reported. An illustrative embodiment of the synergistic combination of treatments leading to the dramatically improved response rate compared to prior methods, is a temperature of about 10°C, mannitol as a carbon source, and 0.05% colchicine.

The inclusion of ascorbic acid, an anti-oxidant, in the isolation medium is preferred for maintaining good microspore viability. However, there seems to be no advantage to including mineral salts in the isolation medium. The osmotic potential of the isolation medium was maintained optimally with about 6% sucrose, although a range of 2% to 12% is within the scope of this invention.

In an embodiment of the embryoid/callus organizing media, mineral salts concentration in IMC Culture Media "D" is (1/8×), the concentration which is used also in anther culture medium. The 6N1 salts major components have been modified to remove ammonium nitrogen. Osmotic potential in the culture medium is maintained with about 12% sucrose and about 400 mg/l proline. Silver nitrate (0.5 mg/l) was included in the medium to modify ethylene activity. The preculture media is further characterized by having a pH of about 5.7 to 6.0. Silver nitrate and vitamins do not appear to be crucial to this medium but do improve the efficiency of the response.

Whole anther cultures can also be used in the production of monocotyledonous plants from a plant culture system. There are some basic similarities of anther culture methods and microspore culture methods with regard to the media used. A difference

from isolated microspore cultures is that undisrupted anthers are cultured, so that a support, e.g., a nylon mesh support, is not needed. The first step in developing the anther cultures is to incubate tassels at a cold temperature. A cold temperature is defined as less than about 25°C. More specifically, the incubation of the tassels is preferably performed at about 10°C. A range of 8 to 14°C is also within the scope of the invention. The anthers are then dissected from the tassels, preferably after surface sterilization using forceps, and placed on solidified medium. An example of such a medium is designated 6N1-TGR-P4.

The anthers are then treated with environmental conditions that are combinations of stresses that are capable of diverting microspores from gametogenesis to embryogenesis. It is believed that the stress effect of sugar alcohols in the preculture medium, for example, mannitol, is produced by inducing starvation at the predefined temperature. In one embodiment, the incubation pretreatment is for about 14 days at 10°C. It was found that treating the anthers in addition with a carbon structure, an illustrative embodiment being a sugar alcohol, preferably mannitol, produces dramatically higher anther culture response rates as measured by the number of eventually regenerated plants, than by treatment with either cold treatment or mannitol alone. These results are particularly surprising in light of teachings that cold is better than mannitol for these purposes, and that warmer temperatures interact with mannitol better.

To incubate the anthers, they are floated on a preculture medium which diverts the microspores from gametogenesis, preferably on a mannitol carbon structure, more specifically, 0.3 M of mannitol plus 50 mg/l of ascorbic acid. Three milliliters is about the total amount in a dish, for example, a tissue culture dish, more specifically, a 60 mm petri dish. Anthers are isolated from about 120 spikelets for one dish yields about 360 anthers.

Chromosome doubling agents can be used in the preculture media for anther cultures. Several techniques for doubling chromosome number (Jensen, 1974; Wan *et al.*, 1989) have been described. Colchicine is one of the doubling agents. However, developmental abnormalities arising from *in vitro* cloning are further enhanced by

colchicine treatments, and previous reports indicated that colchicine is toxic to microspores. The addition of colchicine in increasing concentrations during mannitol pretreatment prior to anther culture and microspore culture has achieved improved percentages.

5 An illustrative embodiment of the combination of a chromosome doubling agent and preculture medium is one which contains colchicine. In a specific embodiment, the colchicine level is preferably about 0.05%. The anthers remain in the mannitol preculture medium with the additives for about 10 days at 10°C. Anthers are then placed on maturation media, for example, that designated 6N1-TGR-P4, for 3 to 6 weeks to induce
10 embryoids. If the plants are to be regenerated from the embryoids, shoot regeneration medium is employed, as in the isolated microspore procedure described in the previous sections. Other regeneration media can be used sequentially to complete regeneration of whole plants.

 The anthers are then exposed to embryoid/callus promoting medium, for example,
15 that designated 6N1-TGR-P4, to obtain callus or embryoids. The embryoids are recognized visually by identification of embryonic-like structures. At this stage, the embryoids are transferred progressively through a series of regeneration media. In an illustrative embodiment, the shoot initiation medium comprises BAP (6-benzyl-amino-
 purine) and NAA (naphthalene acetic acid). Regeneration protocols for isolated
20 microspore cultures and anther cultures are similar.

VIII. ADDITIONAL TISSUE CULTURES AND REGENERATION

 The present invention contemplates a corn plant regenerated from a tissue culture of the inbred maize plant I015036, or of a hybrid maize plant produced by crossing
25 I015036. As is well known in the art, tissue culture of corn can be used for the *in vitro* regeneration of a corn plant. By way of example, a process of tissue culturing and regeneration of corn is described in European Patent Application 0 160 390, the disclosure
1655300.1

of which is incorporated herein by reference. Corn tissue culture procedures are also described in Green and Rhodes (1982) and Duncan *et al.* (1985). The study by Duncan *et al.* (1985) indicates that 97 percent of cultured plants produced calli capable of regenerating plants. Subsequent studies have shown that both inbreds and hybrids
5 produced 91% regenerable calli that produced plants.

Other studies indicate that non-traditional tissues are capable of producing somatic embryogenesis and plant regeneration (Songstad *et al.*, 1988; Rao *et al.*, 1986; Conger *et al.*, 1987; the disclosures of which are incorporated herein by reference). Regenerable cultures, including Type I and Type II cultures, may be initiated from immature embryos
10 using methods described in, for example, PCT Application WO 95/06128, the disclosure of which is incorporated herein by reference in its entirety.

Briefly, by way of example, to regenerate a plant of this invention, cells are selected following growth in culture. Where employed, cultured cells are preferably grown either on solid supports or in the form of liquid suspensions as set forth above. In
15 either instance, nutrients are provided to the cells in the form of media, and environmental conditions are controlled. There are many types of tissue culture media comprising amino acids, salts, sugars, hormones, and vitamins. Most of the media employed to regenerate inbred and hybrid plants have some similar components; the media differ in the composition and proportions of their ingredients depending on the particular application
20 envisioned. For example, various cell types usually grow in more than one type of media, but exhibit different growth rates and different morphologies, depending on the growth media. In some media, cells survive but do not divide. Various types of media suitable for culture of plant cells have been previously described and discussed above.

An exemplary embodiment for culturing recipient corn cells in suspension cultures
25 includes using embryogenic cells in Type II (Armstrong and Green, 1985; Gordon-Kamm *et al.*, 1990) callus, selecting for small (10 to 30 mm) isodiametric, cytoplasmically dense cells, growing the cells in suspension cultures with hormone containing media,

subculturing into a progression of media to facilitate development of shoots and roots, and finally, hardening the plant and readying it metabolically for growth in soil.

Meristematic cells (*i.e.*, plant cells capable of continual cell division and characterized by an undifferentiated cytological appearance, normally found at growing points or tissues in plants such as root tips, stem apices, lateral buds, *etc.*) can be cultured (U.S. Patent No. 5,736,369, the disclosure of which is specifically incorporated herein by reference).

Embryogenic calli are produced essentially as described in PCT Application WO 95/06128. Specifically, inbred plants or plants from hybrids produced from crossing an inbred of the present invention with another inbred are grown to flowering in a greenhouse. Explants from at least one of the following F_1 tissues: the immature tassel tissue, intercalary meristems and leaf bases, apical meristems, immature ears and immature embryos are placed in an initiation medium which contain MS salts, supplemented with thiamine, agar, and sucrose. Cultures are incubated in the dark at about 23°C. All culture manipulations and selections are performed with the aid of a dissecting microscope.

After about 5 to 7 days, cellular outgrowths are observed from the surface of the explants. After about 7 to 21 days, the outgrowths are subcultured by placing them into fresh medium of the same composition. Some of the intact immature embryo explants are placed on fresh medium. Several subcultures later (after about 2 to 3 months) enough material is present from explants for subdivision of these embryogenic calli into two or more pieces.

Callus pieces from different explants are not mixed. After further growth and subculture (about 6 months after embryogenic callus initiation), there are usually between 1 and 100 pieces derived ultimately from each selected explant. During this time of culture expansion, a characteristic embryogenic culture morphology develops as a result of careful selection at each subculture. Any organized structures resembling roots or root primordia are discarded. Material known from experience to lack the capacity for

sustained growth is also discarded (translucent, watery, embryogenic structures). Structures with a firm consistency resembling at least in part the scutelum of the *in vivo* embryo are selected.

5 The callus is maintained on agar-solidified MS or N6-type media. A preferred hormone is 2,4-D. A second preferred hormone is dicamba. Visual selection of embryo-like structures is done to obtain subcultures. Transfer of material other than that displaying embryogenic morphology results in loss of the ability to recover whole plants from the callus.

10 Cell suspensions are prepared from the calli by selecting cell populations that appear homogeneous macroscopically. A portion of the friable, rapidly growing embryogenic calli is inoculated into MS or N6 Medium containing 2,4-D or dicamba. The calli in medium are incubated at about 27°C on a gyrotary shaker in the dark or in the presence of low light. The resultant suspension culture is transferred about once every three to seven days, preferably every three to four days, by taking about 5 to 10 ml of the
15 culture and introducing this inoculum into fresh medium of the composition listed above (PCT Application WO 95/06128).

For regeneration of type I or type II callus, callus is transferred to a solidified culture medium which includes a lower concentration of 2,4-D or other auxins than is present in culture medium used for callus maintenance (PCT Application WO 95/06128,
20 specifically incorporated herein by reference). Other hormones which can be used in regeneration media include dicamba, NAA, ABA, BAP, and 2-NCA. Regeneration of plants is completed by the transfer of mature and germinating embryos to a hormone-free medium, followed by the transfer of developed plantlets to soil and growth to maturity. Plant regeneration is described in PCT Application WO 95/06128.

25 Cells from the meristem or cells fated to contribute to the meristem of a cereal plant embryo at the early proembryo, mid proembryo, late proembryo, transitional or early coleoptilar stage may be cultured so as to produce a proliferation of shoots or multiple

meristems from which fertile plants may be regenerated. Alternatively, cells from the meristem or cells fated to contribute to the meristem of a cereal plant immature ear or tassel may be cultured so as to produce a proliferation of shoots or multiple meristems from which fertile plants may be regenerated (U.S. Patent No. 5,736,369).

5 Progeny of any generation are produced by taking pollen and selfing, backcrossing, or sibling crossing regenerated plants by methods well known to those skilled in the arts. Seeds are collected from the regenerated plants. Alternatively, progeny of any generation may be produced by pollinating a regenerated plant with its own pollen or pollen of a second maize plant. Using the methods described herein, tissue cultures and immature or
10 mature plant tissues may be used as recipient cell cultures for the process of genetic transformation.

IX. PROCESSES OF PREPARING CORN PLANTS AND THE CORN PLANTS PRODUCED BY SUCH CROSSES

15 The present invention also provides a process of preparing a novel corn plant and a corn plant produced by such a process. In accordance with such a process, a first parent corn plant is crossed with a second parent corn plant wherein at least one of the first and second corn plants is the inbred corn plant I015036. An important aspect of this process is that it can be used for the development of novel inbred lines. For example, the inbred
20 corn plant I015036 could be crossed to any second plant, and the resulting hybrid progeny each selfed for about 5 to 7 or more generations, thereby providing a large number of distinct, pure-breeding inbred lines. These inbred lines could then be crossed with other inbred or non-inbred lines and the resulting hybrid progeny analyzed for beneficial characteristics. In this way, novel inbred lines conferring desirable characteristics could be
25 identified.

In selecting a second plant to cross with I015036 for the purpose of developing novel inbred lines, it will typically be desired choose those plants which either themselves

exhibit one or more selected desirable characteristics or which exhibit the desired characteristic(s) when in hybrid combination. Examples of potentially desired characteristics include greater yield, better stalks, better roots, resistance to insecticides, herbicides, pests, and disease, tolerance to heat and drought, reduced time to crop maturity, better agronomic quality, higher nutritional value, and uniformity in germination times, stand establishment, growth rate, maturity, and fruit size. Alternatively, the inbred variety I015036 may be crossed with a second, different inbred plant for the purpose of producing hybrid seed which is sold to farmers for planting in commercial production fields. In this case, a second inbred variety is selected which confers desirable characteristics when in hybrid combination with the first inbred line.

Corn plants (*Zea mays L.*) can be crossed by either natural or mechanical techniques. Natural pollination occurs in corn when wind blows pollen from the tassels to the silks that protrude from the tops of the recipient ears. Mechanical pollination can be effected either by controlling the types of pollen that can blow onto the silks or by pollinating by hand.

In a preferred embodiment, crossing comprises the steps of:

- (a) planting in pollinating proximity seeds of a first and a second parent corn plant, and preferably, seeds of a first inbred corn plant and a second, distinct inbred corn plant;
- (b) cultivating or growing the seeds of the first and second parent corn plants into plants that bear flowers;
- (c) emasculating flowers of either the first or second parent corn plant, *i.e.*, treating the flowers so as to prevent pollen production, or alternatively, using as the female parent a male sterile plant, thereby providing an emasculated parent corn plant;

(d) allowing natural cross-pollination to occur between the first and second parent corn plants;

(e) harvesting seeds produced on the emasculated parent corn plant; and, where desired,

5 (f) growing the harvested seed into a corn plant, preferably, a hybrid corn plant.

Parental plants are typically planted in pollinating proximity to each other by planting the parental plants in alternating rows, in blocks or in any other convenient planting pattern. Where the parental plants differ in timing of sexual maturity, it may be
10 desired to plant the slower maturing plant first, thereby ensuring the availability of pollen from the male parent during the time at which silks on the female parent are receptive to pollen. Plants of both parental parents are cultivated and allowed to grow until the time of flowering. Advantageously, during this growth stage, plants are in general treated with fertilizer and/or other agricultural chemicals as considered appropriate by the grower.

15 At the time of flowering, in the event that plant I015036 is employed as the male parent, the tassels of the other parental plant are removed from all plants employed as the female parental plant to avoid self-pollination. The detasseling can be achieved manually but also can be done by machine, if desired. Alternatively, when the female parent corn plant comprises a cytoplasmic or nuclear gene conferring male sterility, detasseling may
20 not be required. Additionally, a chemical gametocide may be used to sterilize the male flowers of the female plant. In this case, the parent plants used as the male may either not be treated with the chemical agent or may comprise a genetic factor which causes resistance to the emasculating effects of the chemical agent. Gametocides affect processes or cells involved in the development, maturation or release of pollen. Plants treated with
25 such gametocides are rendered male sterile, but typically remain female fertile. The use of chemical gametocides is described, for example, in U.S. Patent No. 4,936,904, the disclosure of which is specifically incorporated herein by reference in its entirety.

Furthermore, the use of Roundup herbicide in combination with glyphosate tolerant maize plants to produce male sterile corn plants is disclosed in U.S. Patent Application Serial No. 08/927,368 and PCT Publication WO 98/44140.

5 Following emasculation, the plants are then typically allowed to continue to grow and natural cross-pollination occurs as a result of the action of wind, which is normal in the pollination of grasses, including corn. As a result of the emasculation of the female parent plant, all the pollen from the male parent plant is available for pollination because tassels, and thereby pollen bearing flowering parts, have been previously removed from all plants of the inbred plant being used as the female in the hybridization. Of course, during
10 this hybridization procedure, the parental varieties are grown such that they are isolated from other corn fields to minimize or prevent any accidental contamination of pollen from foreign sources. These isolation techniques are well within the skill of those skilled in this art.

Both parental inbred plants of corn may be allowed to continue to grow until
15 maturity or the male rows may be destroyed after flowering is complete. Only the ears from the female inbred parental plants are harvested to obtain seeds of a novel F₁ hybrid. The novel F₁ hybrid seed produced can then be planted in a subsequent growing season in commercial fields or, alternatively, advanced in breeding protocols for purposes of developing novel inbred lines.

20 Alternatively, in another embodiment of the invention, both first and second parent corn plants can come from the same inbred corn plant, *i.e.*, from the inbred designated I015036. Thus, any corn plant produced using a process of the present invention and inbred corn plant I015036, is contemplated by the current inventor. As used herein, crossing can mean selfing, backcrossing, crossing to another or the same inbred, crossing
25 to populations, and the like. All corn plants produced using the inbred corn plant I015036 as a parent are, therefore, within the scope of this invention.

The utility of the inbred plant I015036 also extends to crosses with other species. Commonly, suitable species will be of the family *Graminaceae*, and especially of the genera *Zea*, *Tripsacum*, *Coix*, *Schlerachne*, *Polytoca*, *Chionachne*, and *Trilobachne*, of the tribe *Maydeae*. Of these, *Zea* and *Tripsacum*, are most preferred. Potentially suitable
5 for crosses with I015036 can also be the various varieties of grain sorghum, *Sorghum bicolor* (L.) Moench.

A. F₁ Hybrid Corn Plant and Seed Production

Any time the inbred corn plant I015036 is crossed with another, different, corn
10 inbred, a first generation (F₁) corn hybrid plant is produced. As such, an F₁ hybrid corn plant may be produced by crossing I015036 with any second inbred maize plant. Therefore, any F₁ hybrid corn plant or corn seed which is produced with I015036 as a parent is part of the present invention. An example of such an F₁ hybrid which has been produced with I015036 as a parent is the hybrid 8012681. Hybrid 8012681 was produced
15 by crossing inbred corn plant I015036 with the inbred corn plant designated 01INL1 (U.S. Patent No. 6,046,390, the disclosure of which is specifically incorporated herein by reference in its entirety).

The goal of the process of producing an F₁ hybrid is to manipulate the genetic complement of corn to generate new combinations of genes which interact to yield new or
20 improved traits (phenotypic characteristics). A process of producing an F₁ hybrid typically begins with the production of one or more inbred plants. Those plants are produced by repeated crossing of ancestrally related corn plants to try to combine certain genes within the inbred plants.

Corn has a diploid phase which means two conditions of a gene (two alleles)
25 occupy each locus (position on a chromosome). If the alleles are the same at a locus, there is said to be homozygosity. If they are different, there is said to be heterozygosity. In a completely inbred plant, all loci are homozygous. Because many loci when

homozygous are deleterious to the plant, in particular leading to reduced vigor, less kernels, weak and/or poor growth, production of inbred plants is an unpredictable and arduous process. Under some conditions, heterozygous advantage at some loci effectively bars perpetuation of homozygosity.

5 Inbreeding requires sophisticated manipulation by human breeders. Even in the extremely unlikely event inbreeding rather than crossbreeding occurred in natural corn, achievement of complete inbreeding cannot be expected in nature due to well known deleterious effects of homozygosity and the large number of generations the plant would have to breed in isolation. The reason for the breeder to create inbred plants is to have a
10 known reservoir of genes whose gametic transmission is predictable.

 The development of inbred plants generally requires at least about 5 to 7 generations of selfing. Inbred plants are then cross-bred in an attempt to develop improved F₁ hybrids. Hybrids are then screened and evaluated in small scale field trials. Typically, about 10 to 15 phenotypic traits, selected for their potential commercial value,
15 are measured. A selection index of the most commercially important traits is used to help evaluate hybrids. FACT, an acronym for Field Analysis Comparison Trial (strip trials), is an on-farm experimental testing program employed by DEKALB Genetics Corporation to perform the final evaluation of the commercial potential of a product.

 During the next several years, a progressive elimination of hybrids occurs based on
20 more detailed evaluation of their phenotype. Eventually, strip trials (FACT) are conducted to formally compare the experimental hybrids being developed with other hybrids, some of which were previously developed and generally are commercially successful. That is, comparisons of experimental hybrids are made to competitive hybrids to determine if there was any advantage to further development of the experimental
25 hybrids. Examples of such comparisons are presented hereinbelow. After FACT testing is complete, determinations may be made whether commercial development should proceed for a given hybrid.

When the inbred corn plant I015036 is crossed with another inbred plant to yield a hybrid, the original inbred can serve as either the maternal or paternal plant. For many crosses, the outcome is the same regardless of the assigned sex of the parental plants.

However, there is often one of the parental plants that is preferred as the maternal
5 plant because of increased seed yield and production characteristics. Some plants produce tighter ear husks leading to more loss, for example due to rot. There can be delays in silk formation which deleteriously affect timing of the reproductive cycle for a pair of parental inbreds. Seed coat characteristics can be preferable in one plant. Pollen can be shed better by one plant. Other variables can also affect preferred sexual assignment of a
10 particular cross. In the case of the instant inbred, it was generally preferable to use I015036 as the female parent.

B. F₁ Hybrid Comparisons

As mentioned above, hybrids are progressively eliminated following detailed evaluations of their phenotype, including formal comparisons with other commercially
15 successful hybrids. Strip trials are used to compare the phenotypes of hybrids grown in as many environments as possible. They are performed in many environments to assess overall performance of the new hybrids and to select optimum growing conditions. Because the corn is grown in close proximity, environmental factors that affect gene expression, such as moisture, temperature, sunlight, and pests, are minimized. For a
20 decision to be made to commercialize a hybrid, it is not necessary that the hybrid be better than all other hybrids. Rather, significant improvements must be shown in at least some traits that would create improvements in some niches.

Examples of such comparative data are set forth hereinbelow in Table 4, which presents a comparison of performance data for the hybrid 8012681, a hybrid made with
25 I015036 as one parent, versus selected hybrids of commercial value (DK521 and DK527).

All the data in Table 4 represents results across years and locations for research and/or strip trials. The "NTEST" represents the number of paired observations in designated tests at locations around the United States.

1655300.1

TABLE 4: Comparative Data of Hybrid 8012681

HYBRID	NTEST	SI %C	YLD BU	MST PTS	STL %	RTL %	DRP %	FLSTD %M	SV RAT	ELSTD %M	PHT INCH	EHT INCH	BAR %	SG RAT	TST LBS	FGDU	ESTR DAYS
8012681	F 3	113.2	194.7	18.3	2.6	0.0	0.0	96.4	4.7		89.7	41.0		7.0	52.3	1446	98.8
DK521		109.0	194.5	20.6	1.6	0.0	0.0	103.5	5.3		86.7	40.7		5.0	54.1	1510	101.9
DIFF		4.2	0.2	-2.3	1.0	0.0	0.0	-7.1	-0.7		3.0	0.3		2.0	-1.9	-64	-3.1
SIG															*		
8012681	F 1	107.7	172.2	15.6	0.0			97.3	5.0		84.0	42.0		9.0	59.4		101.5
DK527		90.5	155.6	15.9	0.0			105.3	5.0		87.0	42.0		9.0	59.4		101.9
DIFF		17.2	16.6	-0.3	0.0			-8.0	0.0		-3.0	0.0		0.0	0.0		-0.4
SIG																	

Significance levels are indicated as: + = 10%, * = 5%,
** = 1%

LEGEND ABBREVIATIONS:

HYBD = Hybrid

NTEST = Research/FACT

SI %C = Selection Index (percent of check)

YLD BU/A = Yield (bushels/acre)

MST PTS = Moisture

STL % = Stalk Lodging (percent)

RTL % = Root Lodging (percent)

DRP% = Dropped Ears (percent)

FLSTD %M = Final Stand (percent of test mean)

SV RAT = Seedling Vigor Rating

ELSTD %M = Early Stand (percent of test
mean)

PHT INCH = Plant Height (inches)

EHT INCH = Ear Height (inches)

BAR % = Barren Plants (percent)

SG RAT = Staygreen Rating

TST LBS = Test Weight (pounds)

FGDU = GDUs to Shed

ESTR DAYS = Estimated Relative

Maturity (days)

C. Physical Description of F₁ Hybrids

The present invention provides F₁ hybrid corn plants derived from the corn plant I015036. The physical characteristics of an exemplary hybrid produced using I015036 as one inbred parent are set forth in Table 5, which concerns 8012681. An explanation of terms used in Table 5 can be found in the Definitions, set forth hereinabove.

Table 5: Morphological Traits for the 8012681 Phenotype

	CHARACTERISTIC	VALUE
1.	STALK	
	Diameter (width) cm.	2.4
	Anthocyanin	Absent
	Nodes With Brace Roots	1.1
	Brace Root Color	Absent
	Internode Direction	Straight
	Internode Length cm.	16.2
2.	LEAF	
	Color	Green
	Length cm.	82.8
	Width cm.	9.0
	Sheath Anthocyanin	Absent
	Sheath Pubescence	Moderate
	Marginal Waves	Moderate
	Longitudinal Creases	Few
3.	TASSEL	
	Length cm.	44.6
	Spike Length cm.	21.9
	Peduncle Length cm.	13.3
	Branch Number	32.0
	Anther Color	Pink
	Glume Color	Green
	Glume Band	Absent
4.	EAR	
	Silk Color	Green-Yellow
	Number Per Stalk	1.0
	Position (attitude)	Pendant

	Length cm.	17.3
	Shape	Semi-Conical
	Diameter cm.	4.8
	Weight gm.	187.5
	Shank Length cm.	8.5
	Husk Bract	Short
	Husk Cover cm.	1.2
	Husk Opening	Loose
	Husk Color Fresh	Green
	Husk Color Dry	Buff
	Cob Diameter cm.	2.3
	Cob Color	Red
	Shelling Percent	89.7

5.	KERNEL	
	Row Number	18.0
	Number Per Row	36.9
	Row Direction	Straight
	Type	Dent
	Cap Color	Yellow
	Side Color	Yellow
	Length (depth) mm.	12.4
	Width mm.	7.3
	Thickness	3.8
	Weight of 1000K gm.	339.5
	Endosperm Type	Normal
	Endosperm Color	Yellow

**These are typical values. Values may vary due to environment. Other values that are substantially equivalent are also within the scope of the invention. Substantially equivalent refers to quantitative traits that when compared do not show statistical differences of their means.*

5

X. GENETIC COMPLEMENTS

The present invention provides a genetic complement of the inbred corn plant variety designated I015036. Further provided by the invention is a hybrid genetic complement, wherein the complement is formed by the combination of a haploid genetic complement from I015036 and another haploid genetic complement. Means for determining such a genetic complement are well-known in the art.

As used herein, the phrase "genetic complement" means an aggregate of nucleotide sequences, the expression of which defines the phenotype of a corn plant or a cell or tissue of that plant. By way of example, a corn plant is genotyped to determine a representative sample of the inherited markers it possesses. Markers are alleles at a single locus. They
5 are preferably inherited in codominant fashion so that the presence of both alleles at a diploid locus is readily detectable, and they are free of environmental variation, *i.e.*, their heritability is 1. This genotyping is preferably performed on at least one generation of the descendant plant for which the numerical value of the quantitative trait or traits of interest are also determined. The array of single locus genotypes is expressed as a profile of
10 marker alleles, two at each locus. The marker allelic composition of each locus can be either homozygous or heterozygous. Homozygosity is a condition where both alleles at a locus are characterized by the same nucleotide sequence or size of a repeated sequence. Heterozygosity refers to different conditions of the gene at a locus. A preferred type of genetic marker for use with the invention is simple sequence repeats (SSRs), although
15 potentially any other type of genetic marker could be used, for example, restriction fragment length polymorphisms (RFLPs), amplified fragment length polymorphisms (AFLPs), single nucleotide polymorphisms (SNPs), and isozymes.

A genetic marker profile of an inbred may be predictive of the agronomic traits of a hybrid produced using that inbred. For example, if an inbred of known genetic marker
20 profile and phenotype is crossed with a second inbred of known genetic marker profile and phenotype it is possible to predict the phenotype of the F₁ hybrid based on the combined genetic marker profiles of the parent inbreds. Methods for prediction of hybrid performance from genetic marker data is disclosed in U.S. Patent No. 5,492,547, the disclosure of which is specifically incorporated herein by reference in its entirety. Such
25 predictions may be made using any suitable genetic marker, for example, SSRs, RFLPs, AFLPs, SNPs, or isozymes.

SSRs are genetic markers based on polymorphisms in repeated nucleotide sequences, such as microsatellites. A marker system based on SSRs can be highly

informative in linkage analysis relative to other marker systems in that multiple alleles may be present. Another advantage of this type of marker is that, through use of flanking primers, detection of SSRs can be achieved, for example, by the polymerase chain reaction (PCR™), thereby eliminating the need for labor-intensive Southern hybridization. The
5 PCR™ detection is done by use of two oligonucleotide primers flanking the polymorphic segment of repetitive DNA. Repeated cycles of heat denaturation of the DNA followed by annealing of the primers to their complementary sequences at low temperatures, and extension of the annealed primers with DNA polymerase, comprise the major part of the methodology. Following amplification, markers can be scored by gel electrophoresis of
10 the amplification products. Scoring of marker genotype is based on the size (number of base pairs) of the amplified segment.

Means for performing genetic analyses using SSR polymorphisms are well known in the art. The SSR analyses reported herein were conducted by Celera AgGen in Davis, CA. This service is available to the public on a contractual basis. This analysis was
15 carried out by amplification of simple repeats followed by detection of marker genotypes using gel electrophoresis. Markers were scored based on the size of the amplified fragment.

The SSR genetic marker profile of the parental inbreds and exemplary resultant hybrid described herein were determined. Because an inbred is essentially homozygous at
20 all relevant loci, an inbred should, in almost all cases, have only one allele at each locus. In contrast, a diploid genetic marker profile of a hybrid should be the sum of those parents, *e.g.*, if one inbred parent had the allele 168 (base pairs) at a particular locus, and the other inbred parent had 172, the hybrid is 168.172 by inference. Subsequent generations of progeny produced by selection and breeding are expected to be of genotype
25 168, 172, or 168.172 for that locus position. When the F₁ plant is used to produce an inbred, the locus should be either 168 or 172 for that position. Surprisingly, it has been observed that in certain instances, novel SSR genotypes arise during the breeding process. For example, a genotype of 170 may be observed at a particular locus position from the

cross of parental inbreds with 168 and 172 at that locus. Such a novel SSR genotype may further define the parental inbreds from which it was derived. An SSR genetic marker profile of I015036 is presented in Table 6.

5 **Table 6: SSR Profile of Corn Variety I015036 and Comparative Corn Varieties**

LOCUS	I015036	FBLL	FBMU
BNGL105	94	94	96
BNGL118	110	110	110
BNGL149	187	187	187
BNGL244	145	-	145
BNGL426	127	119	127
BNGL589	175	175	175
BNGL615	208	227	209
DUP14	112	112	112
DUP28	123	123	123
MC1014	161	161	161
MC1017	196	196	196
MC1018	130	130	130
MC1022	67	67	67
MC1028	159	159	161
MC1043	175	175	175
MC1046	202	202	-
MC1065	230	230	230
MC1070	220	220	239
MC1074	186	186	174
MC1079	175	177	175
MC1094	178	178	182
MC1108	144	122	144
MC1129	204	204	-
MC1138	186	186	186
MC1176	218	220	218
MC1182	106	106	106
MC1189	219	219	228
MC1191	206	206	203
MC1194	143	143	143
MC1208	111	111	127
MC1209	184	184	184
MC1257	187	187	229
MC1302	149	147	149
MC1305	160	160	160

MC1325	177	177	-
MC1329	93	93	93
MC1360	141	141	145
MC1371	132	124	132
MC1429	191	191	191
MC1456	188	188	188
MC1484	124	124	124
MC1520	275	275	275
MC1523	199	199	199
MC1538	213	237	213
MC1605	128	128	128
MC1662	161	161	161
MC1720	241	241	241
MC1732	100	110	100
MC1740	120	120	120
MC1782	228	228	228
MC1784	250	250	250
MC1808	137	137	137
MC1831	186	184	186
MC1834	216	215	216
MC1839	186	186	-
MC1866	119	119	119
MC1890	136	136	136
MC1904	183	183	183
MC1917	109	109	109
MC1931	174	171	174
MC1940	222	222	216
MC2047	144	144	144
MC2086	242	242	242
MC2122	236	254	236
MC2132	240	240	223
MC2238	195	234	-
MC2259	203	173	203
MC2305	180	190	180
NC004	156	156	156
NC009	147	-	147
PHI017	113	-	113
PHI024	171	171	171
PHI031	194	231	194
PHI033	257	257	257
PHI037	141	137	141
PHI050	92	92	92
PHI051	149	149	149

PHI061	93	85	93
PHI064	104	104	104
PHI065	158	158	158
PHI072	149	149	149
PHI078	129	129	129
PHI089	92	92	92
PHI093	293	293	299
PHI096	109	109	-
PHI101	102	102	102
PHI116	177	-	177
PHI119	171	172	172
PHI120	73	73	73

Primers used to detect SSRs are from Celera AgGen, Inc., 1756 Picasso Ave., Davis, CA 95616

Another aspect of this invention is a plant genetic complement characterized by a genetic isozyme typing profile. Isozymes are forms of proteins that are distinguishable, for example, on starch gel electrophoresis, usually by charge and/or molecular weight. The techniques and nomenclature for isozyme analysis are described in, for example, Stuber *et al.* (1988), which is incorporated by reference.

A standard set of loci can be used as a reference set. Comparative analysis of these loci is used to compare the purity of hybrid seeds, to assess the increased variability in hybrids compared to inbreds, and to determine the identity of seeds, plants, and plant parts. In this respect, an isozyme reference set can be used to develop genotypic "fingerprints."

Table 7 lists the identifying numbers of the alleles at isozyme loci types, and represents the exemplary genetic isozyme typing profile for I015036.

Table 7: Isozyme Profile of Corn Variety I015036 and Comparative Corn Varieties

LOCI	ISOZYME ALLELE		
	I015036	FBLL	FBMU
Acph1	2	2	2
Idh1	4	4	4
Idh2	4	4	4
Mdh1	6	6	6
Mdh2	3.5	3.5	3.5
Mdh3	16	16	16
Mdh4	12	12	12
Mdh5	12	12	12
Pgm1	9	9	9
Pgm2	4	4	4
6Pgd1	3.8	3.8	3.8
6Pgd2	5	5	5

5 The present invention also provides a hybrid genetic complement formed by the combination of a haploid genetic complement of the corn plant I015036 with a haploid genetic complement of a second corn plant. Means for combining a haploid genetic complement from the foregoing inbred with another haploid genetic complement can comprise any method for producing a hybrid plant from I015036. It is contemplated that such a hybrid genetic complement can be prepared using *in vitro* regeneration of a tissue culture of a hybrid plant of this invention.

A hybrid genetic complement contained in the seed of a hybrid derived from I015036 is a further aspect of this invention. An exemplary hybrid genetic complement is that of the hybrid 8012681.

15 Table 8 shows the identifying numbers of the alleles for the hybrid 8012681, which constitutes an exemplary SSR genetic marker profile for hybrids derived from the inbred of the present invention. Table 8 concerns 8012681, which has I015036 as one inbred parent.

Table 8: SSR Profile of Hybrid 8012681

LOCUS	
BNGL105	94.92
BNGL118	110.125
BNGL149	187.183
BNGL426	127.115
BNGL589	175.175
BNGL615	208.231
DUP14	112.103
DUP28	123.133
MC1014	161.161
MC1017	196.196
MC1018	130.138
MC1022	67.116
MC1043	175.189
MC1046	202.206
MC1065	230.241
MC1070	220.254
MC1074	186.164
MC1079	175.173
MC1094	178.170
MC1108	144.144
MC1129	204.204
MC1138	186.190
MC1176	218.220
MC1182	106.82
MC1189	219.227
MC1191	206.207
MC1194	143.143
MC1208	111.111
MC1209	184.184
MC1257	187.179
MC1302	149.147
MC1305	160.200
MC1325	177.171
MC1329	93.93
MC1360	141.143
MC1371	132.134
MC1429	191.191
MC1456	188.176
MC1484	124.124
MC1520	275.275

MC1523	199.199
MC1538	213.213
MC1605	128.110
MC1662	161.161
MC1720	241.241
MC1732	100.100
MC1740	120.167
MC1782	228.228
MC1784	250.254
MC1808	137.131
MC1831	186.184
MC1834	216.208
MC1839	186.194
MC1866	119.123
MC1904	183.191
MC1931	174.170
MC1940	222.216
MC2047	144.144
MC2086	242.247
MC2122	236.236
MC2132	240.223
MC2238	195.195
MC2259	203.180
MC2305	180.190
NC004	156.148
PHI017	113.110
PHI024	171.177
PHI031	194.198
PHI033	257.257
PHI037	141.137
PHI050	92.92
PHI051	149.149
PHI061	93.93
PHI064	104.92
PHI065	158.158
PHI072	149.149
PHI078	129.133
PHI089	92.92
PHI093	293.293
PHI096	109.109
PHI101	102.103
PHI116	177.181
PHI119	171.176

PHI120

73.76

Primers used to detect SSRs are from Celera AgGen, Inc., 1756 Picasso Ave., Davis, CA 95616

The exemplary hybrid genetic complements of hybrid 8012681 may also be assessed by genetic isozyme typing profiles using a standard set of loci as a reference set, using, e.g., the same, or a different, set of loci to those described above. Table 9 lists the identifying numbers of the alleles at isozyme loci types and presents the exemplary genetic isozyme typing profile for the hybrid 8012681, which is an exemplary hybrid derived from the inbred of the present invention. Table 9 concerns 8012681, which has IO15036 as one inbred parent.

Table 9: Isozyme Profile for Hybrid 8012681

Loci	Isozyme Allele
Acph1	2/4
Idh1	4/4
Idh2	4/6
Mdh1	6/6
Mdh2	3.5/3.5
Mdh3	16/16
Mdh4	12/12
Mdh5	12/12
Pgm1	9/9
Pgm2	4/4
6-Pgd1	3.8/2
6-Pgd2	5/5

All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of the foregoing illustrative embodiments, it will be apparent to those of skill in the art that variations, changes, modifications, and alterations may be applied to the composition, methods, and

in the steps or in the sequence of steps of the methods described herein, without departing from the true concept, spirit, and scope of the invention. More specifically, it will be apparent that certain agents that are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope, and concept of the invention as defined by the appended claims.

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

- 5 Anderson, W.P., *Weed Science Principles*, West Publishing Company, 1983.
- Armstrong and Green, "Establishment and maintenance of friable, embryogenic maize callus and the involvement of L-proline," *Planta*, 164:207-214, 1985.
- Conger, Novak, Afza, Erdelsky, "Somatic Embryogenesis from Cultured Leaf Segments of *Zea Mays*," *Plant Cell Reports*, 6:345-347, 1987.
- 10 Duncan *et al.*, "The Production of Callus Capable of Plant Regeneration from Immature Embryos of Numerous *Zea Mays* Genotypes," *Planta*, 165:322-332, 1985.
- Fehr, "Theory and Technique," In: *Principles of Cultivar Development*, 1:360-376, 1987.
- Gaillard *et al.*, "Optimization of Maize Microspore Isolation and Culture Condition for Reliable Plant Regeneration," *Plant Cell Reports*, 10(2):55, 1991.
- 15 Gordon-Kamm *et al.*, "Transformation of Maize Cells and Regeneration of Fertile Transgenic Plants," *The Plant Cell*, 2:603-618, 1990.
- Green and Rhodes, "Plant Regeneration in Tissue Cultures of Maize: Callus Formation from Stem Protoplasts of Corn (*Zea Mays* L.)," In: *Maize for Biological Research*, 367-372, 1982.
- 20 Jensen, "Chromosome Doubling Techniques in Haploids," *Haploids and Higher Plants—Advances and Potentials, Proceedings of the First International Symposium*, 1974.
- Nienhuis *et al.*, "Restriction Fragment Length Polymorphism Analysis of Loci Associated with Insect Resistance in Tomato," *Crop Science*, 27(4):797-803, 1987.
- 25 Omirulleh *et al.*, "Activity of a chimeric promoter with the doubled CaMV 35S enhancer element in protoplast-derived cells and transgenic plants in maize," *Plant Mol. Biol.*, 21(3):415-428, 1993.

- Pace *et al.*, "Anther Culture of Maize and the Visualization of Embryogenic Microspores by Fluorescent Microscopy," *Theoretical and Applied Genetics*, 73:863-869, 1987.
- Poehlman *et al.*, "Breeding Field Crops," 4th Ed., Iowa State University Press, Ames, IA, pp 132-155 and 321-344, 1995.
- Rao *et al.*, "Somatic Embryogenesis in Glume Callus Cultures," *Maize Genetics Cooperation Newsletter*, 60, 1986.
- Songstad *et al.*, "Effect of 1-Aminocyclopropane-1-Carboxylic Acid, Silver Nitrate, and Norbornadiene on Plant Regeneration from Maize Callus Cultures," *Plant Cell Reports*, 7:262-265, 1988.
- Sprague and Dudley (eds.), "Corn and Corn Improvement," 3rd Ed., Crop Science of America, Inc., and Soil Science of America, Inc., Madison Wisconsin. pp 881-883 and pp 901-918, 1988.
- Stuber *et al.*, "Techniques and scoring procedures for starch gel electrophoresis of enzymes of maize *C. Zea mays*, L.," *Tech. Bull.*, 286, 1988.
- Wan *et al.*, "Efficient Production of Doubled Haploid Plants Through Colchicine Treatment of Anther-Derived Maize Callus," *Theoretical and Applied Genetics*, 77:889-892, 1989.
- Wang *et al.*, "Large-Scale Identification, Mapping, and Genotyping of Single-Nucleotide Polymorphisms in the Human Genome," *Science*, 280:1077-1082, 1998.
- Williams *et al.*, "Oligonucleotide Primers of Arbitrary Sequence Amplify DNA Polymorphisms which Are Useful as Genetic Markers," *Nucleic Acids Res.*, 18:6531-6535, 1990.